

Universal Definition and Classification of Heart Failure

A Report of the Heart Failure Society of America, Heart Failure Association of the European Society of Cardiology, Japanese Heart Failure Society and Writing Committee of the Universal Definition of Heart Failure Consensus Conference

Endorsed by Canadian Heart Failure Society, Heart Failure Association of India, Cardiac
Society of Australia and New Zealand, Chinese Heart Failure Association

WRITING COMMITTEE MEMBERS*

Biykem Bozkurt, MD, PhD, *Chair*
Andrew Coats, MD, Co-Chair
Hiroyuki Tsutsui, MD, Co-Chair

Magdy Abdelhamid, MD
Stamatis Adamopoulos, MD
Nancy Albert, PhD, CCNS, CHF, CCRN, NE-BC
Stefan Anker MD, PhD
John Atherton, MBBS, PhD
Michael Böhm, MD
Javed Butler, MD, MPH, MBA
Mark H. Drazner, MD, MSc
G. Michael Felker, MD, MHS,
Gerasimos Filippatos, MD
Gregg C. Fonarow, MD
Mona Fiuzat, PharmD
Juan –Esteban Gomez-Mesa, MD
Paul Heidenreich, MD
Teruhiko Imamura, MD, PhD
James Januzzi, MD
Ewa A. Jankowska, MD PhD
Prateeti Khazanie, MD, MPH

Koichiro Kinugawa, MD, PhD
Carolyn S.P. Lam, MBBS, FRCP, PhD
Yuya Matsue, MD, PhD
Marco Metra, MD
Tomohito Ohtani, MD, PhD
Massimo Francesco Piepoli, MD, PhD
Piotr Ponikowski, MD, PhD
Giuseppe M.C. Rosano, MD, PhD
Yasushi Sakata, MD, PhD
Petar Seferović, MD, PhD

Randall C. Starling, MD, MPH
John R. Teerlink, MD
Orly Vardeny, PharmD, MS
Kazuhiro Yamamoto, MD, PhD
Clyde Yancy, MD, MSc,
Jian Zhang, MD, PhD
Shelley Zieroth, MD

Address for Correspondence:

Biykem Bozkurt, MD, PhD, FHFSA, FACC, FAHA
The Mary and Gordon Cain Chair of Cardiology and Professor of Medicine
Medical Care Line Executive, DeBakey VA Medical Center
Director, Winters Center for Heart Failure Research
Associate Director, Cardiovascular Research Institute
Associate Provost of Faculty Affairs and Senior Associate Dean of Faculty Development
Baylor College of Medicine
2002 Holcombe Blvd, Houston, Texas 77030 Email: bbozkurt@bcm.edu
Phone: 713-794-7350, Fax: 713-79407551, Twitter handle: @BiykemB

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ejhf.2115

*The Comprehensive RWI Data Supplement table is available with this article in Appendix 1. Affiliations are also listed in Appendix 1.

This article is co-published in *Journal of Cardiac Failure* and *European Journal of Heart Failure*

Table of Contents

| | |
|-------------------------------------------------------------------------------------------------------|----|
| Preamble..... | 4 |
| 1. Methodology..... | 4 |
| 1.1 Writing Committee Composition | 5 |
| 1.2. Consensus Development | 5 |
| 1.3. Peer Review and Approval..... | 5 |
| 2. Current Definitions of HF | 6 |
| 2.1. Definitions of HF Used in Current Clinical Trials and Registries | 6 |
| 2.2 Gaps in Current Definitions of HF | 8 |
| 3. Current Classifications of Heart Failure | 12 |
| 3.1. Current Classification of Heart Failure According to Ejection Fraction and Its Limitations | 12 |
| 3.2 Current Classification According to Stages of Heart Failure and Its Limitations | 14 |
| 4. Gaps in Definitions According to Trajectory of Changes in HF | 17 |
| 4.1. Trajectory Changes in EF..... | 17 |
| 4.2 Trajectory Changes in Clinical Status..... | 17 |
| 5. Learning from Other Disease Definitions | 5 |
| 6. Proposed Universal Definition of Heart Failure..... | 20 |
| 7. Other Syndromes Related to Heart Failur | 23 |
| 8. Proposed New Stages of Heart Failure | 28 |
| 9. Proposed New Classifications of HF According to Ejection Fraction..... | 32 |
| 10. Approaches to Specific Etiology of HF | 33 |
| References..... | 36 |
| Tables | 51 |
| Figures..... | 53 |
| Appendix 1. Author Relationships With Industry and Other Entities | 57 |
| Appendix 2. Abbreviations | 69 |
| | |

Summary of Key Points

- In this document, we propose a universal definition of heart failure (HF) as the following: HF is a clinical syndrome with symptoms and or signs caused by a structural and/or functional cardiac abnormality and corroborated by elevated natriuretic peptide levels and or objective evidence of pulmonary or systemic congestion.
- We propose revised stages of HF as: At-risk for HF (Stage A), for patients at risk for HF but without current or prior symptoms or signs of HF and without structural or biomarkers evidence of heart disease. Pre-heart failure (Stage B) for patients without current or prior symptoms or signs of HF but evidence of structural heart disease or abnormal cardiac function, or elevated natriuretic peptide levels. HF (Stage C) for patients with current or prior symptoms and/or signs of HF caused by a structural and/or functional cardiac abnormality. Advanced HF (Stage D) for patients with severe symptoms and/ or signs of HF at rest, recurrent hospitalizations despite guideline-directed management and therapy (GDMT), refractory or intolerant to GDMT, requiring advanced therapies such as consideration for transplant, mechanical circulatory support, or palliative care.
- Finally, we propose a new and revised classification of HF according to left ventricular ejection fraction (LVEF). The classification includes HF with reduced EF (HFrEF): HF with $LVEF \leq 40\%$; HF with mid-range EF (HFmrEF): HF with $LVEF 41-49\%$; HF with preserved EF (HFpEF): HF with $LVEF \geq 50\%$; and HF with improved EF (HFimpEF): HF with a baseline $LVEF \leq 40\%$, a ≥ 10 point increase from baseline LVEF, and a second measurement of $LVEF > 40\%$

Preamble

Currently available definitions of heart failure (HF) are ambiguous and lack standardization.¹⁻⁸ Some definitions focus on the diagnostic features of the clinical syndrome,³⁻⁵ whereas other definitions approach the definition as a characterization of the hemodynamic and physiological aspects.^{2, 8} There is significant variation in different platforms,¹⁻⁵ and a growing need for standardization of the definition of HF.^{6, 9}

A Universal Definition of HF is of critical importance to clinicians, investigators, administrators, health care services, institutions, legislators and payers alike. The increasing prevalence and burden of HF^{10, 11}; an increased recognition of growing health care disparities¹² and deficiencies in the optimal treatment in HF with guideline directed medical treatment (GDMT) strategies^{13, 14} all underline the necessity for a universally recognizable definition of HF. Evolving evidence for new effective preventive and treatment strategies in HF will require clarity in the different stages and/or ejection fraction (EF) subgroups of HF,^{15, 16} along with an increased emphasis on performance measures with a need for accuracy in patient diagnoses and treatment indications^{17,18, 19}; a need for improved communication and understanding of the definition of HF with patients and for shared decision making and transitions of care between different levels of care and health care professionals³ and an increased recognition and emphasis of standard diagnoses and endpoints in the settings of research, clinical trials and registries^{20, 21}.

The objectives of this document are to provide a universal definition of HF that is clinically relevant, simple but conceptually comprehensive, with the ability to sub-classify and to encompass stages within; with universal applicability globally, and with prognostic and therapeutic validity and acceptable sensitivity and specificity. We envision the proposed universal definition and classifications to be used in a standardized fashion across scientific societies and guidelines, employed by clinicians, and used in research studies.

1. Methodology

1.1 Writing Committee Composition

The Heart Failure Society of America (HFSA), Heart Failure Association of the European Society of Cardiology (HFA) and the Japanese Heart Failure Society selected the members of the writing committee. The writing committee consisted of 37 individuals with domain expertise in HF, cardiomyopathy and cardiovascular disease.

1.2. Consensus Development

On August 20, 2020, in response to the necessity for consensus for definition of HF, the Heart Failure Society of America (HFSA), Heart Failure Association of the European Society of Cardiology (HFA) and the Japanese Heart Failure Society convened a virtual consensus

conference to develop a universal definition of heart failure with participation from fourteen different countries and six continents. The work of the writing committee was accomplished via a series of teleconference and Web conference meetings, along with extensive email correspondence. The review work was distributed among subgroups of the writing committee based on interest and expertise. The proceedings of the workgroups were then assembled, resulting in the proposed universal definition. All members reviewed and approved the final vocabulary.

1.3. Peer Review and Approval

The 2020 Universal Definition of HF was reviewed by official reviewers nominated by the HFSA, HFA and JHFS. The writing committee anticipates that the proposed definition and classification will require review and updating in the same manner as other published universal definitions.²² The writing committee, therefore, plans to review the universal definition on a periodic basis, starting with the anniversary of publication of the definition, to ascertain whether modifications should be considered.

2. Current Definitions of Heart Failure

HF is a clinical syndrome with different etiologies and pathophysiology rather than a specific disease. This makes defining HF more complex than diseases that have a pathologic gold standard for diagnosis such as cancer. Not surprisingly, definitions of HF vary widely in the medical literature, in contemporary guidelines, and in medical practice. Differing definitions have been developed for different purposes, ranging from “textbook” definitions of HF, which are typically focused on pathophysiology, to case definitions such as the Framingham criteria²³ that are primarily used in research. The traditional textbook definition of HF, which is usually defined as a “condition in which the heart cannot pump enough blood to meet the body's needs”¹ or “abnormality of cardiac structure or function leading to failure of the heart to deliver oxygen at a rate commensurate with the requirements of the metabolizing tissues”,² is a complex and impractical definition that often cannot be verified in practice and apply to only a certain subgroup of patients with HF. As such, in a study of advanced HF patients awaiting left ventricular assist device implantation, cardiac output was shown to be insufficient to meet the metabolic needs of the body only in 25 % of these advanced HF patients at rest, demonstrating

the inadequacy of such definitions in the majority of the HF population.²⁴ In clinical care, other diagnostic criteria such as measurement of plasma natriuretic peptides play an important role in clarifying the diagnosis of HF.³⁻⁵ A summary of contemporary definitions of HF from the American College of Cardiology/American Heart Association (ACC/AHA); Heart Failure Association/European Society of Cardiology (HFA/ESC), and JHFS guidelines is provided in **Table 1**. Although the definitions of HF used in current practice guidelines from the ACC/AHA³, HFA/ESC⁴, and JHFS⁵ differ in some details, they share the following common elements: they identify HF as a *clinical syndrome*, i.e., a recognizable cluster of signs and symptoms; they require the presence of at least some of the cardinal symptoms of HF including dyspnea, fluid retention/edema, fatigue, activity intolerance and exercise limitation; and they require some form of structural or functional heart disease. Some also specify a reduced cardiac output and/or elevated intra-cardiac pressures at rest or during stress.⁴ Overall, the existing definitions of HF comprise three elements: evidence of structural heart disease, a history of symptoms that are commonly reported in heart failure and objective signs commonly seen in HF.

2.1. Definitions of Heart Failure Used in Current Clinical Trials and Registries

The definitions and inclusion criteria used in clinical trials and registries in HF differ from those in clinical practice, guidelines and textbooks. Most trials in HF with reduced left ventricular ejection fraction (HFrEF) (**Table 2**), and in HF with preserved left ventricular ejection fraction (HFpEF) (**Table 3**) reflect inclusion criteria that usually include a LVEF threshold, an established HF diagnosis with specific NYHA Class categories, certain levels of natriuretic peptides and may sometimes include a requirement of past HF hospitalizations, depending on the severity of HF targeted for the trial. HFpEF studies also may include corroborative evidence by imaging reflecting structural and or functional changes. Nonetheless, a number of gaps remain in standardizing criteria for clinical trials. These include sensitivity and specificity of diagnostic criteria for HF, establishing standardized natriuretic peptide criteria; the complexity of additional requirements to ascertain the diagnosis of HF; challenges with HFpEF including multiple comorbidities which are often excluded in clinical trials, how to handle patients with EF recovery or changes in clinical trajectory, competing diagnoses that may mimic findings of HF and generalizability of the trial criteria to the ultimately intended treatment

population. It is also important to distinguish between clinical trial inclusion criteria that aim to select target populations, from clinical trial end-point definitions that facilitate measurement of outcomes secondary to the disease process. For example, natriuretic peptides, which are commonly used in entry criteria in HF trials, are not commonly required for clinical end-point definitions.²¹

2.2 Gaps in Current Definitions of HF

Combined Definition with Hemodynamic Characterization of Heart Failure

The current definitions that include a hemodynamic characterization such as the HFA/ESC definition which defines HF as a “a clinical syndrome characterized by typical signs and symptoms, caused by a structural and/or functional cardiac abnormality, resulting in a reduced cardiac output and/or elevated intra-cardiac pressures at rest or during stress”,⁴ have the following limitations: while accurate, this type of definition is hard to apply in public health or epidemiological settings, because of the subjectivity of the symptoms counterbalanced by the unfeasibility (invasive) or unreliability of measurements of cardiac output or filling pressures. For a definition to be also useful for the non-specialist it should be assessable easily and with relatively low inter-observer variability. Framingham criteria, which were developed for just such a purpose²³ are now considered insufficiently specific for adoption as a definition of HF in the contemporary setting.

Cardiomyopathy and Heart Failure

A key distinction that has led to persistent confusion in many discussions of the definition of HF is that between the concepts of “heart failure” and “cardiomyopathy.” As defined above, HF is a clinical syndrome, i.e., a recognizable pattern of signs and symptoms. “Cardiomyopathy” is a term, itself with widely differing definitions, that describes features of structural and functional heart muscle dysfunction. These different definitions may lead to potential confusion. In clinical practice, the term “cardiomyopathy” is often used as a more general term encompassing types of cardiac dysfunction, that may be further qualified with the underlying cause (e.g., ischemic cardiomyopathy, non-ischemic cardiomyopathy, etc.). Alternatively, cardiomyopathy may be understood to be a specific form of myocardial disease that excludes forms of HF with a clearly established cause (such as ischemic heart disease). Even guideline statements from various scientific bodies have varied in their use of this term.^{25, 26} Furthermore, the maladaptive

hemodynamic and compensatory mechanisms in HF may result in development of or worsening of cardiomyopathy.²⁷ Classification systems have been proposed that attempt to incorporate both the classification of HF and cardiomyopathy into a unified system, most notably the proposed MOGES criteria (Morpho-functional phenotype-M, organ(s) involvement -O, genetic inheritance pattern-G, etiological annotation -E including genetic defect or underlying disease/substrate, and the functional status -S), but these have not been widely adopted due to their complexity.²⁸ HF encompasses a broader spectrum of cardiac disorders, not only cardiomyopathies that could be an underlying cause of the HF syndrome. In this statement, we do not provide specific classification strategies for cardiomyopathies, which we believe to be outside the scope of this document.²⁵

Biomarkers in the Definition of Heart Failure

Natriuretic peptides such as BNP and NT-proBNP are elevated in most forms of HF and are an integral component of making a diagnosis of HF in many clinical settings, especially when the diagnosis is uncertain.^{3-5, 29} Use of these biomarkers has the highest class of recommendation to support a diagnosis or exclusion of HF^{4, 29} in contemporary practice guidelines, but are notably absent from most definitions of HF. This is in contrast to the universal definition of myocardial infarction (MI), where elevations of a circulating biomarker (troponin) are both central to the clinical diagnosis and fundamental to the universal definition itself.²² Although a biomarker based approach has incremental diagnostic value, especially in the context of clinical uncertainty, in some communities with limited resources, natriuretic peptide measurements currently may not be readily available, but their availability is rapidly increasing, and natriuretic peptide measurements are becoming part of standard care. Furthermore, certain clinical conditions other than HF, such as chronic kidney disease, atrial fibrillation, pericardial disease, pulmonary embolism, and even aging can also result in an increase in natriuretic peptide levels, and obesity is associated with lower natriuretic peptide levels, underlining the importance of an individualized interpretation of biomarker levels, particularly in special populations and in the setting of competing diagnoses and comorbidities. It is important to recognize that though measuring natriuretic peptide levels may improve diagnostic accuracy and guide risk stratification in patients with HF, in certain patients with HF, such as patients with HFpEF or obesity, natriuretic peptide levels can be lower than those without HFrEF (though usually higher

than those without HF); this may complicate their use for diagnosis and prognosis.. Differences according to race/ethnicity, sex and age will need to be taken into consideration in interpretation and different thresholds are commonly used for patients with atrial fibrillation, a very common comorbidity in HF that can lead to increased natriuretic peptide levels. A potential influence of comorbidities is also relevant for troponin interpretation in patients with suspected acute MI, but despite similar limitations, the introduction of a quantitative biomarker element to a disease definition has improved the accurate classification of disease states and proven to be of value in MI and other diseases.^{22,30} In general, both BNP and NT-proBNP values track similarly, and either can be used in patient care settings as long as their respective absolute values and cut points are not used interchangeably. Notably, BNP, but not NT-proBNP, is a substrate for neprilysin. Angiotensin receptor neprilysin inhibitor (ARNi) may result in an increase in BNP levels, but not NT-proBNP levels.²⁹ Furthermore, patient level changes need to be interpreted according to baseline levels; natriuretic peptides are higher during periods of decompensation compared to compensated periods, reflecting dynamic temporal changes.

Clinical and Research Aspects of Defining Heart Failure

Clinical research requires standardized definitions for identifying cases of HF and the collection of endpoints of interest, including especially HF-related hospitalizations (HFH).²¹ Given the increased use of electronic health records (EHR) as research tools, there is growing interest in the use of computer algorithms to identify cases of HF from EHR data for research purposes. Although classical signs and symptoms are often included in EHR data they may not be codified as discrete data fields, leading to increased interest in the use of machine learning techniques to identify cases.³¹ Definitions of HF are important not only for clinical practice or research entry criteria, but also for generalizability of research findings to the HF population, uniformity in endpoints of clinical trials; reliability and appropriateness of data captured in clinical, administrative and billing registries and performance measures.

Patient and Clinician Perspective

A syndrome that is based solely on symptoms can be confusing for clinicians and patients, both because they are often not specific to a single disease (e.g. fatigue and dyspnea) and because they are highly subjective, for example with the same objective limitation being considered disabling by one person and perceived as being normal for age by another. Once diagnosed, and with

effective therapy, patients may become asymptomatic (NYHA Class I); however, structural, cellular and molecular abnormalities may continue to worsen silently.³² Although Stage C HF uses the wording “current or previous symptoms” in the definition, patients may believe that lack of signs and symptoms equates to “being out of HF”, and be less likely to adhere to care.³³ Health care professionals may be less likely to optimize GDMT when symptoms are mild or absent.³⁴ Removing the word “congestive” in the term HF was an important reminder to providers that there is a range of signs and symptoms once diagnosed. Further, patients may not understand or recognize when HF worsens, until symptoms are severe enough to prompt emergency care.³⁵ In the era of shared decision making and patient understanding of chronic conditions, it will be important to acknowledge and incorporate different stages that are understandable by patients after diagnosis.

Competing diagnoses

There are many conditions that may mimic HF, either in isolation (mimicry) or when co-existing with HF (co-causative). The combination of acute dyspnea, hypervolemia and cardiorenal syndrome is often labeled as HF in an emergency care setting, although the problem could be confounded by, or even be predominantly due to, anemia and iron deficiency. Recognizing proportionate contributions of a clinical picture, to dissect out the element that is specifically HF-related, will be an important part of establishing a HF diagnosis, and it may not be an easy differentiation to make in all situations. It is HF only if the cardiac component is considered “important”. However, it is also important to recognize that HF can co-exist with other diagnoses. For example, HF syndromes with lesser degrees of systolic impairment such as HF with preserved ejection fraction (HFpEF) frequently present with a wide range of cardiac and non-cardiac abnormalities.³⁶ Newer, sometimes inconsistent terminology regarding mid-range ejection fraction has further complicated subcategorization of HF. It is important to promote greater clarity and specificity in the diagnosis of HF.

3. Current Classifications of Heart Failure

An important part of defining HF is that of creating a “usable” classification scheme. There are a variety of classification frameworks in current use that attempt to define distinct subsets of HF (**Table 4**). Some of these, such as New York Heart Association (NYHA) class and EF categories, have been subsequently used as entry criteria in clinical trials, resulting in their

incorporation into product labeling and guideline recommendations about which patients should receive a given therapy.³⁻⁵ Others, such as classifying patients by HF etiology may have important implications for prognosis or differential response to therapy.³⁷

3.1 Current Sub classification of Heart Failure according to Ejection Fraction and Its Limitations

Due to the fact that clinical trial inclusion criteria, and hence evidence of benefit, has often been restricted to patients with a reduced EF, HF has traditionally been subcategorized according to EF when defining recommended treatments in clinical practice guidelines.³⁻⁵ All guidelines use the terminology of HF with reduced EF (HFrEF), and HF with preserved EF (HFpEF) (**Table 5**) but differ in the terminology used in patients with EF's between 40 and 49%. The 2013 ACC/AHA guidelines have used the terminology of HFpEF-borderline for patients with EF between 41-49%, and HFpEF-improved for those whose EF improved from a lower level to EF>40% under the subgrouping of patients with HFpEF.³ The HFA/ESC and JHFS guidelines have defined a third category of HF with mid-range EF (HFmrEF) or mildly reduced EF for those with EF 41-49%.^{4, 5} The concept of HFmrEF is not necessarily accepted by all guidelines.³⁸

In an effort, through a public-private partnership with the US Food and Drug administration (FDA) and with an intent to standardize terminology and LVEF cut-points used in US clinical trials, Heart Failure Collaboratory, and Academic Research Consortium proposed the following definitions and EF ranges as their most recent recommendations: HF with reduced EF (HFrEF): HF with left ventricular ejection fraction (LVEF) $\leq 40\%$. HF with preserved EF: HF with LVEF $\geq 50\%$. HF with mid-range EF (HFmrEF): HF with LVEF $> 40\%$ and LVEF $< 50\%$.³⁹

The dichotomization of left ventricular ejection fraction (LVEF) of above or below e.g. 40% has been helpful to apply therapies that have been shown to work in patients with reduced EF. Further classification into HF mid-range EF (HFmrEF) has potential utility as well as challenges due to its ambiguity, uncertainty and dynamic trajectory.^{15, 40} Post-hoc analyses of certain HF trials have suggested that standard therapy for HFrEF may be effective and extended to patients with HFmrEF,⁴¹⁻⁴⁴ but meta-analyses report diverse findings with neurohormonal antagonism in patients with HFmrEF, specifying benefit in certain subgroups and underlining heterogeneity of

this category^{15, 44-46} The characteristics of HFmrEF overlap with HFrEF and HFpEF; straddling either category; sometimes one more than the other depending on the clinical circumstance or patients studied.¹⁵ In population-based studies, usually without exclusions of specific etiologies, HFmrEF comprises 10-20 % of the HF population;^{43, 47} resembles the HFrEF group, but with similar⁴⁶ or better survival than HFrEF patients.^{47,15} Although some patients' characteristics of HFmrEF are just between those of HFrEF and HFpEF, the prognosis of patients is not necessarily related to EF,⁴⁸ and the relation between mortality and B-type natriuretic peptide (BNP) is not affected by EF.^{48, 49} In many patients, HFmrEF reflects a transitional trajectory for a dynamic temporal change; either to improvement or recovery from HFrEF^{46, 50}, or to deterioration to HFrEF.^{15, 46, 50, 51} Although HFrEF and HFpEF have different clinical spectrums and proposed pathophysiological mechanisms, there is no clear defining syndrome recognized or postulated for HFmrEF. It is likely that patients in this range may have etiologies that are similar to those in lower or higher LVEF groups, and may be in transition from higher to lower LVEF or vice versa. Persistent HFmrEF can be seen in some, including heterogeneous etiologies such as those with ischemic, infiltrative, restrictive or hypertrophic cardiomyopathies.^{46, 50, 51} Therefore, lower than a normal EF does not necessarily represent one phenotype and does not always entail the maladaptive deleterious mechanisms seen in patients with HFrEF. Furthermore, patients with restrictive, infiltrative and hypertrophic cardiomyopathies, who may have HFmrEF, have traditionally been excluded from some clinical trials, emphasizing the necessity to focus on etiology rather than LVEF. The prevalence of HFmrEF, without overlap of other categories, has posed a major challenge for recruitment in trials, resulting in termination due to enrolment futility⁵² and in some clinical trials and epidemiological studies, patients with LVEF 40-49% have been categorized as HFpEF.

Another criticism is the accuracy of the measurement of EF in clinical practice.

Echocardiography is widely used to assess EF in patients with cardiovascular diseases, but the inter-observer and intra-observer variability are not small enough to allow precise quantification of differences in one integer place values such as 39 versus 41%. Although other cardiovascular imaging modalities can be used to assess EF, there is substantial variation between modalities as well.⁵³ Furthermore, EF is not a reliable measure of contractile performance, is load dependent, and can vary according to hemodynamic status and loading conditions. Other imaging modalities

such as global longitudinal strain are evolving to better characterize the ventricle, structural abnormalities, contractile performance, reverse remodeling, response to therapy and will likely expand the structural phenotyping beyond EF.

Finally, the trajectory of EF over time in addition to a single absolute value of EF, and severity of LV dysfunction even among HFrEF may need to be taken into account to further classify patients with HF. Despite all these reservations, classification by EF has proven to be clinically and epidemiologically useful.

3.2 Current Classification According to Stages of Heart Failure and Its Limitations

The ACC/AHA stages are categorized as Stage A, Patients at high risk for HF but without structural heart disease or symptoms of HF; Stage B, Structural heart disease but without signs or symptoms of HF; Stage C, Structural heart disease with prior or current symptoms of HF; Stage D, Refractory HF requiring specialized interventions.^{3, 4, 54} The original ACC/AHA definition of stages of HF⁵⁴ has been ubiquitously adapted throughout other HF guidelines globally.³⁻⁵

Although these stages of HF are well recognized amongst healthcare professionals, they are not standard nomenclature for general practitioners, patients, payers or among the literature or education platforms provided by patient advocacy groups. Patients living with HF are less likely to identify with stages of HF in comparison to the familiarity with EF and subjective symptom burden. Contemporary clinical trials have not enrolled or randomized based on stages of HF and most treatment strategies are not guided by the stages in HF.

The ACC/AHA stages are based on symptoms and the presence or absence of structural heart disease and are applicable to both HFrEF and HFpEF. Certainly, there are prognostic nuances that are missed in such a broad staging classification, and opinions also vary as to whether those individuals solely identified with risk factors should be labeled as having a disease state, especially given that they are risk factors for many different diseases (not just HF risk factors). In comparison classification schemas such as the SCAI cardiogenic shock stages⁵⁵ classified their stages based on detailed parameters of laboratory values, hemodynamics as well as physical exam and exemplifies a more detailed approach to staging. Furthermore, the definitional

progression along the ACC/AHA stages A-D is a unidirectional path with little appreciation of a possibility to revert to a lower stage with appropriate guideline directed medical therapy.

If the HF process were to be defined as a continuum from stage A through D, the highest number of patients would be in stage A or Stage B.⁵⁶⁻⁵⁹ This is due to the fact that the prevalence of hypertension, diabetes, coronary artery disease, obesity/metabolic syndrome- the risk factors with significant relative risk and population attributable risk for development of HF- are present in approximately one third of the US population.¹⁰ By population based registries, more than 40-50% of the adult population have been categorized to be in Stages A or B.⁵⁶⁻⁵⁸ The high prevalence of HF risk in the general population raises the question of whether Stage A patients should really be defined to have HF. From the public and health care perspective, being called HF, regardless of such an early status as stage A, raises important concerns since HF is usually perceived as an advanced chronic disease with symptoms and very adverse outcomes, and may have implications for health and life insurance. Of course, it is critical to focus on prevention, with recognition, prevention and treatment of these risk factors, but it is also important to differentiate those who have HF from those at-risk for HF. Similarly, clinicians in general or HF practice have not adopted the terminology of Stage A HF beyond academic circles, partly due to lack of actionable specific treatment recommendations according to stages, and most of their assessment and management focuses on management of left ventricular dysfunction (Stage B) or symptomatic HF (stages C/D). When clinicians address risk factors such as hypertension, diabetes, obesity or coronary artery disease, they do not refer to those as Stage A HF or pre-HF but rather independent diagnoses. Furthermore, despite recognized increased adverse outcome risk and possibility of progress to symptomatic HF in some^{56, 59, 60}, the data on likelihood of progression from Stages A/B to C/D are limited.^{57, 59, 60} Thus most clinicians do not commonly use the HF terminology for stage A patients, and do not commonly educate patients regarding risk of progression from Stages A/B to C.

Another important development that needs to be taken in consideration of stages in HF is the advances in prevention of future risk of HF by specific therapies. Though in the past, prevention and holistic treatment of risk factors by standard treatment strategies were felt to prevent HF³, there is growing evidence that certain treatment strategies are better for the prevention of HF, and not all treatment strategies of hypertension and diabetes prevent HF equally or at all. For

example, in the treatment of hypertension, diuretic-based antihypertensive therapies, angiotensin converting enzyme (ACE)-inhibitors or angiotensin receptor blockers (ARBs) have been shown to prevent HF in a wide range of target populations, whereas calcium channel blockers have not.⁶¹ There is growing evidence that treatment with sodium–glucose cotransporter 2 (SGLT2)-inhibitors prevent HF hospitalizations among patients with Type II diabetes⁶²⁻⁶⁴ or in patients with HFrEF regardless of diabetes^{65, 66} whereas other glucose treatment strategies do not. It is also interesting to note that patients with a higher future HF risk identified by risk scores that include biomarkers such as albuminuria, appear to confer greater benefit from SGLT2-inhibitor therapy among patients with type II diabetes.⁶⁷ The biomarker profile may identify patients with cardiometabolic, cardiovascular and cardiac structural changes in patients predestined to develop HF or in other words pre-HF. Supporting this concept was the STOP-HF trial which provided evidence that screening with natriuretic peptides among individuals with cardiovascular disease or with cardiovascular risk factors such as diabetes and hypertension, can be helpful to prevent development of HF or left ventricular systolic or diastolic dysfunction.⁶⁸ Accordingly, the 2017 ACC/AHA /HFSa focused update for the management of HF guidelines incorporated recommendations for natriuretic peptides-based screening in the prevention of HF as a Class IIa recommendation.²⁹ Similarly, high sensitivity cardiac troponin levels are associated with future development of incident HF in the general population^{69, 70} and in those with evidence of cardiotoxicity or cardiac injury in high risk populations⁷¹ allowing for treatment strategies to prevent development of HF. Thus, biomarker elaboration can further identify risk and presence of ultrastructural abnormalities in HF among asymptomatic patients and could be a marker for Stage B HF without development of macroscopic structural changes detectable by imaging or ECG.

4. Gaps in Definitions According to Trajectory of Changes in HF

The HF syndrome is dynamic, with changing clinical trajectories based on signs, symptoms, and disease progression, driven by underlying pathophysiologic processes. Changes in HF may be captured in several ways, including alterations in cardiac structure and function and by clinical status.

4.1 Trajectory Changes in EF

GDMT can result in improvement in LVEF and reverse remodeling in patients with HFrEF.⁷² The phenomenon of improvement and recovery of LVEF has led to a growing interest in the long-term outcomes and management of these patients and how they differ from “non-responders”, or individuals whose LVEF does not improve with treatment. Currently, there is no consensus definition for patients with HFrEF whose LVEF improves, which has led to a variety of terms describing this phenotype, including patients with “improved” LVEF, HFpEF (borderline), HFpEF, and HF with recovered EF (HFrecEF). The magnitude of change that defines “recovery” of LVEF is not standardized, but it is recognized that distinguishing HFrecEF from HF with midrange (HFmrEF) requires serial measurements of LVEF to appropriately capture change over time as this group might represent HFrecEF or deteriorated HFpEF. Moreover, since measurement of LVEF is subject to significant intra- or inter-reader variability, small changes in LVEF need to be interpreted cautiously. Thus, a recent scientific panel put forth a working definition of HFrecEF that includes: a baseline LVEF $\leq 40\%$, a $\geq 10\%$ increase from baseline LVEF, and a second measurement of LVEF $> 40\%$.⁷² In this formulation, recovered EF signifies improvement of LVEF to over 40% but not necessarily totally normalized. There have been other attempts to characterize improvement in EF as an increase in LVEF by more than 10%.⁷² It is also important to recognize that the trajectory might not be linear and unidirectional, a patient may have improvement followed by a decline in EF or vice versa depending on underlying etiology, duration of disease, adherence to the GDMT, comorbidities or re-exposure to cardiotoxins.

4.2 Trajectory Changes in Clinical Status

Another method that captures the HF trajectory relies on an assessment of the patient’s clinical status, which can inform the risk for hospitalization for HF or for mortality. A *de novo* diagnosis of HF, also referred to as *new onset HF*, carries an increased risk for adverse clinical outcomes since the patient is not likely to be treated with optimal GDMT at the time of diagnosis.

Most patients with HF have episodes of clinical *worsening of HF*, which has been previously defined as worsening signs or symptoms in concert with a hospitalization.⁷³ Data from more contemporary studies resulted in expansion of worsening HF to also include patients who require escalation of outpatient therapies, such as diuretics, even without a hospitalization.⁷⁴ This is

because the need for intensifying diuretic therapy, regardless of location (inpatient or outpatient), portends a worse prognosis than a patient who does not require intensification of therapy.

Worsening HF implies a period of stability preceding a deterioration of signs and symptoms.

However, the phrase “stable” HF may be a misnomer, as patients with HF always carry a residual risk for hospitalization or sudden cardiac death, even when minimally symptomatic or asymptomatic receiving optimal treatment. For such patients *remission* may be a more suitable term.⁷⁵ When a patient with worsening HF does not improve with therapy escalation and continues to decline, s/he can be referred to as *refractory* to treatment. These patients are often assessed for advanced therapies such as mechanical circulatory support or cardiac transplant, or if they do not qualify for advanced therapies, clinicians consider referral for palliative care.

Patients may have *improvement* in HF symptoms, functional capacity, quality of life and exercise performance with GDMT. Some patients with reversible or treatable causes of HF such as cardiomyopathy due to hypertensive heart disease, alcoholic cardiomyopathy, peripartum cardiomyopathy, or tachycardia-induced cardiomyopathy may even *recover* from HF with treatment and manifest resolution of HF symptoms, normalization of EF and cardiac structure. These patients require close follow-up and require continuation of treatment to ascertain that HF symptoms or LV dysfunction do not reoccur in the future.⁷⁶

5. Learning from Other Disease Definitions

Disease definitions are not all the same. Some are categorical, where the disease is present or it is not. In some there may be a single pathognomonic feature that defines the disease state, such as many cancers and infectious diseases. In others, where numerical thresholds are used, a disease may be defined against a quantitative threshold of abnormality in an anatomical and/or functional feature. Examples of these include hypertension, osteoporosis, sarcopenia and chronic kidney disease (CKD). In some (CKD, hypertension) the presence of this numerical abnormality alone is sufficient to define the disease, while in others (e.g. HF, sarcopenia) the loss of function must be symptomatic or functionally evident for the disease to be defined. In the current Universal Definition of Myocardial Infarction (MI), elevation of cardiac troponin is central to the clinical diagnosis and fundamental to the universal definition.²²

There are many other corollaries and lessons to learn from other areas of cardiology and medicine in regards to disease definition and classifications. Current ACC/AHA classification of valvular heart disease is very similar to the current ACC/AHA HF categorization into Stages A-D.^{3, 77} Such categorization is an epidemiology-based system where the disease stage is defined based on stages of susceptibility from ‘at risk’ to subclinical disease to clinical disease, and finally, recovery, disability, or death. Atrial fibrillation is also based on an epidemiology-based system where patients are categorized as paroxysmal (≤ 48 hours), persistent (>7 d or cardioverted), long-standing (>1 yr), and permanent.⁷⁸ However, in atrial fibrillation, clinicians also use the CHA2DS2-VASc risk score to determine potential stroke risk and thereby guide management.⁷⁸ A similar parallel in HF is the MAGGIC model for prediction of mortality and other attempts at scoring to help risk stratify patients who may have worsening HF, rehospitalization, or higher chance of dying.⁷⁹

In regards to non-cardiovascular strategies for disease definition, there are quite a few examples. CKD is classified based on albuminuria and eGFR.³⁰ Albuminuria states are similar to numeric categorization of disease, like LVEF in HF, while eGFR ranging from normal to end-stage renal disease provides prognostic information and guides management decisions, such as drug dosing and need for dialysis. Liver disease is categorized based on pathology using imaging and tissue sampling to define levels of steatosis, hepatitis, fibrosis, and cirrhosis.⁸⁰ Much like CKD, liver disease also supplements disease categorization with risk scores like the MELD score.⁸¹ Lung disease is assessed using pulmonary function tests which helps clinicians stratify patients based on air-flow limitation and the GOLD system.⁸² Chronic obstructive lung disease also stratifies patients based on symptoms and risk of exacerbations similar to congestion-perfusion⁸³ categorization in HF. Pulmonary hypertension classification (WHO Groups 1-5)⁸⁴ is similar to the etiology-based groupings for cardiomyopathies^{25, 26} with genetic, acquired, and mixed categories and is a potential model for future HFpEF⁸⁵ disease stratification. Finally, the field of cancer groups disease using a combination of epidemiology-based staging (i.e., at risk for cancer, pre-cancer, carcinoma in situ, localized, early/late locally advanced, and metastasized) coupled with disease-specific markers that determine treatment course and targeted therapies. Cancer, which is a chronic disease similar to HF, reflects one of the most comprehensive combined approaches of classification using epidemiology, biomarker thresholds and trajectory.

Future attempts at defining HF will need to draw on principles of categorization used in other disease states. Each organ system has unique pathophysiology that helps determine its disease categorization, and ultimately, all organ systems are interconnected. Indeed, HF represents an end-stage phenotype for most (if not all) cardiovascular diseases. In the terminal stages of disease, the universal element is disseminated disease and multiorgan failure. However, unlike other organ systems, the heart is unique in that hemodynamics play a central role in the disease state. Many disease states are moving towards a **combination** of epidemiology-based, numeric, and targeted marker-based therapies. Disease definitions are critical to patients' and clinicians' understanding of their pathology, informs clinical decision making, categorization for financial billing, and creation of future health policies.

6. Proposed Universal Definition of Heart Failure

In this section, we provide a consensus opinion on a new proposed universal definition of HF.

Symptoms

HF, like many non-categorical diseases, is widely held to be a clinical syndrome, devoid of any single pathognomonic histological or biochemical signal, and being the possible end result of many quite distinct and varied clinical disease states. Common symptoms and signs of HF are listed in **Table 6**.

The current ACCF/AHA Classification of HF³ includes two pre-symptomatic stages, A and B. Though we restrict the *definition* of syndrome of HF to being a symptomatic clinical condition, our proposed revised stages still straddle the pre-symptomatic stages. To not lose the advantage that the A/B/C/D staging system offered, to incorporate the asymptomatic phases under the HF umbrella, and to enhance understandability of these asymptomatic phases we propose a new categorization of Stages A and B into “at risk” and “pre-HF” in section 9 below.

Objective Marker

In learning from other disease states that incorporated a core and frequently measured variables in their definition, such as acute MI, eGFR in CKD, HBA1_c in diabetes, bone mineral density in osteoporosis or FEV₁ in COPD, making the diagnosis more accessible to non-specialists and

more reliable and consistent between observers, hospitals and health care systems, we propose incorporation of an objective measurement in addition to the symptoms in the HF definition.

In HF, possible candidates for such a measurement might theoretically be hemodynamic measures such as elevated pulmonary capillary wedge pressure and right atrial pressure by right heart catheterization, biomarkers associated with congestion such as natriuretic peptides, measures of neurohormonal over activity or measures of exercise limitation such as maximal oxygen consumption. None of these are commonly or reliably associated with the disease states of HF, e.g. LVEF can vary from low through normal to high and still be part of an HF syndrome; no single hemodynamic measure is adequate to serve as a practical, non-invasive and reliable measurement; measurement of exercise limitation with cardiopulmonary exercise testing with expired gas exchange is not practical or universally available; and to date, neurohormone levels have not universally been considered reliable measures of the disease state. The closest have been the natriuretic peptides (NP's), which are recommended in modern guidelines as both diagnostic tests of reasonable clinical usefulness with prognostic utility and as good tests to rule out HF as a cause of breathlessness in certain settings.^{4, 29} Contemporary guidelines already state that NPs can be used as an initial diagnostic test, and that patients with normal plasma NP concentrations are unlikely to have HF.^{4, 29} A detailed diagnostic algorithm will require specific operational thresholds based on individual NP's and assay systems, as well as detailing other clinical features which can affect NP levels, but for common clinical purposes simple thresholds can be established which have sufficient operational accuracy to be incorporated usefully into a universal definition of HF.

Proposed New HF definition

We propose a contemporary universal definition of HF (**Figure 1**) that is simple but conceptually comprehensive, with near universal applicability, prognostic and therapeutic validity, and acceptable sensitivity and specificity.

❖ UNIVERSAL HEART FAILURE DEFINITION

HF is a clinical syndrome with current or prior

- **Symptoms and or signs (Table 6) caused by a structural and/or functional cardiac abnormality (as determined by EF<50%, abnormal cardiac chamber**

enlargement, E/E' >15, moderate/severe ventricular hypertrophy or moderate/severe valvular obstructive or regurgitant lesion)

- **and corroborated by *at least one* of the following:**
 - elevated natriuretic peptide levels (for values refer to table 7)
 - objective evidence of cardiogenic pulmonary or systemic congestion by diagnostic modalities such as imaging (e.g. by CXR or elevated filling pressures by echocardiography) or hemodynamic measurement (e.g. right heart catheterization, PA catheter) at rest or with provocation (e.g. exercise)

Such a definition is comprehensive and practical enough to form the base which allows further sub-classifications and which can encompass formal disease stages, both with universal applicability, prognostic and therapeutic validity, and an acceptable sensitivity and specificity. Please note that the definition of HF requires not only symptoms or signs, (**Table 6**) but presence of either elevated natriuretic peptides or objective evidence of pulmonary or systemic congestion by diagnostic modalities. For example, it would be important for peripheral edema or ascites (Table 6) to be corroborated by presence of elevated right-sided cardiac filling pressures or rales by presence of elevated left sided cardiac filling pressures; or elevated natriuretic peptides. It is also important to note that elevated jugular venous pressure estimate by an experienced clinician could be accepted as an objective evidence.

Please also note that in certain patients, congestion and hemodynamic abnormalities may become manifest with provocation such as exercise, especially in patients with HFpEF. This can support the diagnosis of HF. It is also critical note that in patients with low perfusion and hypovolemic state, there may not be any evidence of congestion or elevated filling pressures, but rather decreased cardiac output accompanied with low or normal ventricular filling pressures,⁸⁶ (e.g. in the setting of over-diuresis in patients with HF). Once hypovolemic state is corrected, patients with HF usually have elevated filling pressures.

In the definition above, we did not specify left or right HF. Though left heart HF, and in advanced stages, biventricular HF are common, right HF can also be recognized as part of the

above definition when patients present with symptoms or signs (Table 6) caused by a cardiac abnormality and have elevated natriuretic peptide levels or objective evidence of cardiogenic pulmonary or systemic congestion. Right HF primarily due to cardiac abnormalities such as arrhythmogenic right ventricular cardiomyopathy (ARVC) would be part of this definition.

We recognize that asymptomatic stages with patients at-risk (former Stage A HF), with structural heart disease or cardiomyopathies (former Stage B HF) would not be covered under the above definition as having HF, which emphasizes symptoms and signs of HF, but we conceptualize the HF syndrome as a continuum of disease with certain stages as pre-HF. This is similar to the approach with other disease states such as cancer, which defines those at risk and pre-cancer. The stages preceding the symptomatic phases as those **at -risk** and **pre-heart failure** will be discussed in the following section.

We also realize certain patients with competing diagnoses such as CKD with marked volume overload, can present with symptoms and signs of HF, have elevated natriuretic peptides, and may even have evidence of congestion by imaging or elevated filling pressures. Though some of these patients may have concomitant HF, these patients have a primary abnormality that may require a specific treatment beyond HF. In the following section, we will address such other syndromes.

7. Other Syndromes Related to Heart Failure

As noted above, the definition of HF comprises a combination of symptoms and/or signs of HF caused by a structural and/or functional cardiac abnormality, and evidence of elevated filling pressures by natriuretic peptides or by imaging/ hemodynamic assessment. Although many clinicians will initially envision patients with left HF as embodying this definition, it is important to note that there are other syndromes that may fulfill this definition of HF, addressed below. These etiologies require specific treatment and management strategies targeting the underlying or proximate cause, as well as treating HF itself.

Right Heart Failure

The most common cause of right HF is left HF. However, right HF is characterized not only by signs and symptoms of right sided HF but also by right atrial enlargement or right ventricular

dysfunction. The presence of right HF in the setting of left HF is typically due to post-capillary, WHO group 2 pulmonary hypertension and may require modified treatment approaches and portends a poor prognosis; therefore, recognition of biventricular HF is important.⁸⁴ Given the importance of these distinctions, the classification of types of ventricular failure in HF commonly includes three categories; left ventricular failure (LVF), right ventricular failure (RVF) and combined left and right ventricular failure usually termed as biventricular failure. We believe isolated right HF due to primary pulmonary hypertension etiologies (WHO Groups 1, 3, 4), though may have symptoms or signs which may mimic HF and may have elevated natriuretic peptide levels, would not be categorized under HF, as the signs and/or symptoms are not *caused* primarily by a structural and/or functional cardiac abnormality. On the other hand, right HF due to primary right ventricular conditions such as arrhythmogenic right ventricular cardiomyopathy (ARVC) would be categorized under HF.

Acute Myocardial Infarction /Acute coronary syndrome

Acute myocardial infarction may be complicated by HF. Given its acuity, specific pathophysiology and specific treatment strategies; we believe acute myocardial infarction would be the overarching definition for the episode in proximity to acute myocardial infarction. It is also possible that these patients may recover with timely treatment strategies and not progress to chronic HF, but also many may progress to chronic HF. In clinical trials, patients with acute myocardial infarction or acute coronary syndrome within six weeks are usually excluded from clinical trials in HF. These patients may present with asymptomatic left ventricular dysfunction or pre-HF or symptoms and signs of HF due to a cardiac abnormality and may have elevated natriuretic peptides or evidence of congestion by imaging or hemodynamics. During the acute phase, these patients are diagnosed as myocardial infarction complicated by HF rather than HF alone. This does not mean acute myocardial infarction should be replaced by HF alone, but it does mean the setting and specific etiology of HF can be an important feature that determines specific therapeutic approaches. This setting has also been subject to specific clinical trial evaluation.⁸⁷⁻⁸⁹ In addition to specific therapies for acute myocardial infarction, these patients have indications for specific treatment for asymptomatic LV dysfunction (pre-HF or Stage B HF) or symptomatic HF complicating acute myocardial infarction during the acute phase, or as primary diagnoses in the chronic phase post myocardial infarction.

Cardiogenic Shock

Another important form of HF is cardiogenic shock which is the clinical state of organ hypoperfusion due to severe cardiac dysfunction. In cardiogenic shock, the symptoms and signs reflecting HF include hypotension unresponsive to volume repletion, altered mental status, cool extremities, and laboratory evidence of end organ dysfunction such as elevated lactate levels due to hypoperfusion.⁵⁵ Cardiogenic shock is an extreme form of HF which requires some form of definitive therapy such as intravenous inotropes, vasopressors or mechanical circulatory support. Cardiogenic shock is a type of HF, but due to its specific hemodynamic and clinical characterization requiring specific therapies such as vasoactive agents, circulatory support and/or revascularization depending on etiology, we believe keeping the descriptor “cardiogenic shock” will help identify a patient cohort with specific and urgent treatment needs. Cardiogenic shock may occur as an acute de novo presentation (e.g. large acute myocardial infarct, fulminant myocarditis) or with progressive deterioration in a patient with chronic HF. Subacute cardiogenic shock may be in continuum of the wet and cold advanced HF patient with low cardiac output state. Such patients may meet the criteria for cardiogenic shock especially when they have evidence of end-organ dysfunction. A system describing stages of cardiogenic shock has been proposed by SCAI and other societies and characterizes the patients as Stage A “at risk” for cardiogenic shock, stage B “beginning” shock, stage C “classic” cardiogenic shock, stage D “deteriorating”, and E “extremis”.⁵⁵ Such classification is important to characterize the severity and stage of shock, but it is also important to acknowledge the presence of HF as the preceding cause of shock in such patients, and identify advanced HF complicated with cardiogenic shock as the diagnosis.

Hypertensive Emergency and Hypertensive Heart Disease

Hypertensive emergencies encompass a spectrum of clinical presentations of uncontrolled blood pressure associated with end-organ damage that can include acute left ventricular dysfunction, pulmonary edema, myocardial infarction/ischemia, and/or aortic dissection. All of these complications may result in or be complicated with an acute presentation of HF. Hypertension increases HF risk by two to three fold⁹⁰ and accounts for almost half of the HF cases in the US population as a population attributable risk.⁹¹ Thus both acutely hypertensive emergency and chronically, hypertensive heart disease can be complicated with HF. Treatment of hypertension

is upmost importance in prevention and treatment of HF, underlined as a Class I recommendation with strong level of evidence in guidelines.^{4, 29}

Valvular Heart Disease

Aortic stenosis and mitral regurgitation can result in HF. Valvular heart disease is acknowledged as a specific disease, as it results in specific hemodynamic and ventricular alterations and requires specific treatment strategies targeting valvular abnormality. Most HF clinical trials exclude significant valvular heart disease for these reasons.

Congenital Heart Disease

Some types of congenital heart disease can result in HF. Incomplete or palliative correction of a congenital lesion leading to a chronic state of hemodynamic stress may result in subsequent HF, especially in complex congenital heart diseases such as tetralogy of Fallot, single ventricle defects and transposition of the great arteries. Additional myocardial, coronary or conduction system injury can occur due to complications of corrective surgery and can lead to progressive contractile dysfunction in some patients. The treatment should target the underlying anomaly and specific hemodynamic conditions.

High-output Failure

High output HF presents with similar symptoms and signs of systemic or pulmonary congestion, frequently associated with rapid heart rate and signs of peripheral vasodilation. Cardiac dysfunction may be represented by pathologically elevated cardiac output, echocardiographic signs of right ventricular dilation or dysfunction, and elevated natriuretic peptide concentrations. High output HF is a response to extracardiac causes including liver disease, arteriovenous shunt, lung disease, thiamine deficiency, anemia, thyroid disease or myeloproliferative disorders. Treatment is generally directed to the underlying causes. Given the unique nature of high output failure, it is appropriate that it have a separate classification.

Other Overlapping and Competing Diagnoses with HF

Patients can experience clinical deterioration as specific events that may not necessarily meet the universal definition of a diagnosis of HF. Such occurrences consist of events of a primary disease process that may be associated with signs and symptoms of HF as a result of the *primary* cause that is not HF at that encounter. These can include cardiovascular causes such as acute

myocardial infarction or acute coronary syndrome, hypertensive emergency as mentioned above, and also other CV primary diagnoses such as atrial fibrillation with rapid ventricular response, prolonged ventricular arrhythmias, pulmonary embolus, pericardial diseases, and acute valvular dysfunction. In these cardiovascular diagnoses, complication *with* HF is associated with worse prognosis and outcomes and underlines the urgency of addressing the underlining problem as well as the HF.

Other non-cardiovascular entities such as renal failure, liver failure, morbid obesity with peripheral edema and chronic respiratory failure hypoventilation syndrome may present with symptoms and signs that mimic HF. Due to volume overload and neurohormonal compensatory mechanisms involved in some of these disease states, symptoms, signs and even hemodynamic characterization and biomarker profile can overlap with HF, and these patients may indeed also have concomitant HF. In these cases, the *proximate cause* of the signs and symptoms of volume overload is a distinct entity to which treatment is often primarily directed, in addition to HF. These events are often of significant interest to clinical events committees of clinical trials, where they may be considered as an event “*with* HF” rather than a primary HF event. Another important concept that supports the principality of these competing diagnoses are that the symptoms and signs of HF may disappear once the underlying primary cause is treated, for example symptoms and signs that mimic HF may resolve with hemodialysis in a patient with end stage CKD who may have missed a dialysis appointment. Thus, it is important not to catalog every presentation with shortness of breath and edema that requires treatment with fluid management strategies or diuretics as HF. It is however, also important to not miss the complication *with* HF which requires timely management of HF as well as the proximate cause. Many of these factors can contribute to worsening outcomes in a complementary fashion in patients with HF. For example, patients with HF associated with CKD or DM are at much higher risk than those without. Rather than “competing”, these diagnoses can become complementary comorbidity risk factors to HF for worse outcomes.

8. Proposed Revised Stages of the Heart Failure Continuum

To enhance clinician, patient and public understanding and adoption; to avoid the stigma of HF before the symptoms are manifest; to address the evolving role of biomarkers to define patients

with structural and subclinical heart disease who are at higher risk of developing HF and are potential candidates for targeted treatment strategies for the prevention of HF; and to address some of the gaps identified in Section 3.2 in the current approach to staging HF, we propose the following stages. (**Figure 2**)

- ❖ **AT RISK FOR HEART FAILURE (STAGE A):** Patients at risk for HF but without current or prior symptoms or signs of HF and without structural cardiac changes or elevated biomarkers of heart disease. Patients with hypertension, atherosclerotic cardiovascular disease, diabetes, obesity, known exposure to cardiotoxins, positive family history of cardiomyopathy or genetic cardiomyopathy would be in this category. Not all of these patients will develop HF, but risk factor intervention may be warranted.
- ❖ **PRE-HEART FAILURE (STAGE B):** Patients without current or prior symptoms or signs of HF *but* evidence of *one* of the following:
 - Structural Heart Disease: e.g. left ventricular hypertrophy, cardiac chamber enlargement, ventricular wall motion abnormality, myocardial tissue abnormality (e.g. evidence of myocardial edema, scar/fibrosis abnormality by CMR T2 or LGE Imaging), valvular heart disease
 - Abnormal cardiac function: e.g. reduced LV or RV ventricular systolic function, evidence of increased filling pressures (by invasive or noninvasive measures), abnormal diastolic dysfunction
 - Elevated natriuretic peptide levels (for levels, refer to **Table 7**) or elevated cardiac troponin levels (over 99th percentile in a normal reference population) especially in the setting of exposure to cardiotoxins.
- ❖ **HEART FAILURE (STAGE C):** Patients with current or prior symptoms and/ or signs of HF caused by a structural and/or functional cardiac abnormality.
- ❖ **ADVANCED HEART FAILURE (STAGE D) :** Severe symptoms and/ or signs of HF at rest, recurrent hospitalizations despite GDMT, refractory or intolerant to GDMT, requiring advanced therapies such as consideration for transplant, mechanical circulatory support, or palliative care

Abnormal cardiac function: e.g. reduced LV or RV ventricular systolic function, can be characterized by reduced ejection fraction, abnormal ventricular strain, or other noninvasive or invasive modalities.

Though certain genetic markers may be associated with structural cardiac changes and future HF, we did not specifically include genetic markers in the definition of pre-HF or Stage B HF as the penetrance, expressivity, phenotypic characterization and prognosis with genetic markers vary significantly. As evidence for precision for risk evolves with biomarkers, genetics, omics and/or risk calculators, alternative approaches can be developed in the future to identify risk categories beyond traditional risk factors, and pre-HF beyond cardiac structure and biomarkers alone; and support expansion of indications for preventive treatment strategies for patients at risk or with pre-HF.

Please note that the cutoffs provided for natriuretic peptide levels in Table 7 represent thresholds lower than inclusion criteria used in some clinical trials for symptomatic HF,^{65, 66} but similar to those used in former guidelines.⁴ Thresholds proposed in the table have higher sensitivity and may have lower specificity especially in older patients, or patients with atrial fibrillation or chronic kidney disease. Usually, higher cutoff values are recommended for the diagnosis of HF in these patients.⁹² For example, for ages 50-75, NT-proBNP threshold value of 900; for ages >75 years, NT-proBNP value of 1,800 pg/ml provide reliable sensitivity and specificity for the diagnosis of HF, compared to a NT-proBNP value of 450 pg/ml for ages <50 among patients requiring hospitalization.⁹² Similarly, in patients with atrial fibrillation, an increase by 20-30 % have been suggested in natriuretic peptide level thresholds for trial enrollment in HF,⁹² since atrial fibrillation is known to result in increased concentrations of natriuretic peptides even in the absence of HF. Furthermore, it is important to note that natriuretic peptide cut-offs selected for population screening for pre-HF (Stage B HF) may be lower than 99% reference limits⁶⁸ and will need to be defined according to the population at risk.

NYHA Classification

The NYHA classification is important to characterize symptoms and functional capacity of patients with symptomatic (Stage C) HF or advanced HF (Stage D). The NYHA classification system categorizes HF on a scale of I to IV; Class I: No limitation of physical activity, Class II:

Slight limitation of physical activity, Class III: Marked limitation of physical activity, Class IV: Symptoms occur even at rest; discomfort with any physical activity. We believe, it is important to specify NYHA class at baseline after the initial diagnosis, and after treatment through the continuum of care of a patient with HF. A patient with symptomatic HF (Stage C) may become asymptomatic with treatment. Since that patient will still be categorized as HF /Stage C, NYHA Class I can further specify his/her absence of current symptoms. Worsening NYHA Class is associated with worse prognosis and any symptomatic patient with HF (NYHA Class II-IV HF) should have further optimization of GDMT.

Recognition of Clinical Trajectory in Heart Failure

It is well-recognized that the natural history of HF encompasses changes in the clinical risk of hospitalization and death over time, with risk increasing from “pre-HF” to “new onset/ de novo HF”, and further increasing with each episode of “worsening HF” where there is deterioration of HF signs and symptoms despite ongoing therapy, requiring hospitalization or outpatient escalation of therapy.⁹³ It is crucial to identify both the stage of the patient’s natural history, as well as recognize the patient’s clinical trajectory (improving vs stalled or persistent vs worsening),⁹⁴ for optimal treatment, risk mitigation strategies, and patient-centered discussions. Gaining perspective of not only where the patient stands at the point in time, but in which direction the patient is headed, is a critical element of determining whether to continue along the current therapeutic course or to change direction. Thus a patient with “worsening chronic HF” following initial stabilization of “new onset/ de novo HF” would alert a physician of the immediate high risk for recurrent hospitalization or death, particularly in the period of close proximity to the worsening HF event, and trigger escalation of disease modifying therapies rather than a focus on decongestion with diuretics alone. Of note, we caution against a terminology of “stable HF”, as patients are expected to improve with GDMT. These patients should have further optimization of therapies despite perceived stability or improvement, as there is evidence for significant improvement in outcomes with additional therapies in these patients. Lack of improvement is a marker of worse prognosis and should be termed as “**persistent**” rather than “stable”, and prompt clinicians to further optimize therapy. For those patients who have resolution of symptoms and signs of HF along with resolution of previously present structural and functional heart disease after a phase of symptomatic HF, we recommend “**HF in**

remission” or NYHA Class I HF status rather than “recovered HF” which should be reserved for patients who have persistent resolution of HF symptoms and signs, normalization of cardiac structure, function and biomarker profile following resolution and treatment of a *fully reversible* cause, especially in view of the TRED-HF trial results which demonstrated that many patient deemed to have “recovered” from dilated cardiomyopathy will relapse following treatment withdrawal suggesting remission rather than recovery.⁷⁶ (Figure 2) Full and persistent recovery is rare, and even in the setting of reversible causes, patients may have recurrence of symptoms and or develop LV dysfunction in the future.

Acute versus Decompensated Heart Failure

In this document, we do not use the terms “acute new onset HF” or “acute decompensated HF” which are the terminologies commonly used to describe patients requiring hospitalization or urgent care. The indications for hospitalization and or urgent care utilization vary, and most patients who require hospitalization for HF may have chronic progressively worsening HF, rather than an acute singular event. We realize these patients may present with rapid onset or progressively escalating symptoms and/or signs of HF which are associated with adverse outcomes, requiring urgent evaluation and treatment. We have elected to characterize these patients as “decompensated HF” which may represent acutely decompensated patients due to an inciting event (e.g. atrial fibrillation with rapid ventricular response) or chronically and progressively worsening patients with marked deterioration of HF signs and symptoms despite ongoing therapy requiring urgent intervention, hospitalization, or rapid escalation of therapies including advanced therapies.

We recognize there are a variety of acute presentations of HF (e.g. myocarditis, peripartum, cardiotoxicity, stress cardiomyopathy etc.) and other entities associated with acute presentations of HF such as hypertensive emergency and acute myocardial infarction which will require specialized treatment strategies targeting the underlying etiology. These have been addressed by others⁹⁵⁻⁹⁷ and are beyond the scope of this document.

9. Proposed New Classifications of HF According to Ejection Fraction

The strongest argument to use LVEF to categorize HF is that LVEF defines a group known to respond to life-prolonging therapy from randomized controlled trials.^{3, 4, 29, 41, 42, 75, 87, 89, 98-101}

While LVEF also provides prognostic information, this reason alone does not justify using LVEF to define HF. Accordingly, LVEF categories were created that define groups where treatment differs.

To be able to differentiate patients who benefit from GDMT according to clinical trial entry criteria of patients with HFrEF, capture evolving recognition of the need to identify effective treatment strategies in patients with HF associated with a mildly reduced or mid-range LVEF, as well as preserved LVEF, and harmonize with existing practice guidelines, we propose the following four classifications of EF (**Figure 3**):

- ❖ **HF with reduced EF (HFrEF):** HF with $LVEF \leq 40\%$.
- ❖ **HF with mid-range EF (HFmrEF):** HF with $LVEF 41-49\%$
- ❖ **HF with preserved EF (HFpEF):** HF with $LVEF \geq 50\%$.
- ❖ **HF with improved EF (HFimpEF):** HF with a baseline $LVEF \leq 40\%$, a ≥ 10 point increase from baseline LVEF, and a second measurement of $LVEF > 40\%$

We acknowledge the growing body of evidence that standard therapy for HFrEF may be effective in and extended to select patients with HFmrEF.⁴¹⁻⁴⁴ It is however important to recognize the heterogeneity of this category, underlined by diverse findings from meta-analyses with neurohormonal antagonism, specifying benefit in certain subgroups.^{15, 44-46}

Evidently, LVEF is not a singular measurement by which LV function is assessed in isolation. Chamber volumes and other cardiac structural and functional parameters are important and other diagnostic modalities can be complementary. Though the above classification is provided for targeting GDMT according to LVEF indications, other cardiac features are also important for phenotypic characterization, etiology or prognosis. Development of LV dilation in a patient with HFpEF or HFmrEF may imply impending HFrEF. It is important to recognize that cardiac structural and functional information in addition to LVEF is important to guide management of the patient.

Since GDMT can result in improvement in LVEF and reverse remodeling in patients with HFrEF, the trajectory of improvement and recovery of EF has been of interest to determine the types (e.g. device, medical, advanced) and duration of treatment.⁷² In cases where longitudinal surveillance of LVEF is available, clinicians should also consider the trajectory of the LVEF, in addition to the LVEF at the point in time, recognizing that a significant reduction in LVEF over time is a poor prognostic factor calling for consideration of intensification of therapy and advanced management strategies according to patient goals. Importantly, EF can decline after withdrawal of pharmacological treatment in many patients who had improved EF to normal range with GDMT.⁷⁶ This implies that there is not full recovery in cardiac structure and function in most patients despite improvement in EF. Therefore, we recommend use of the improved terminology rather than recovered EF. We believe “improved EF” deserves a separate classification and should not be classified as HFmrEF or HFpEF even after improvement in LVEF to 41-49% or $\geq 50\%$ respectively, as discontinuing HFrEF therapy in this group leads to poor outcome.⁷⁶ GDMT should be continued in patients with HFimpEF regardless of whether it has improved to normal range, LVEF $\geq 50\%$, especially in view of the TRED-HF trial results.⁷⁶ We also recognize that patients with baseline LVEF of 41-49 % who have improved LVEF to $\geq 50\%$ may be categorized as HFimpEF.

10. Approaches to Specific Etiologies of HF

In addition to the recognition of the syndrome of HF and its classifications, it is critical that every effort should be made to diagnose and define the specific etiology/etiologies of HF. Understanding the underlying etiological processes of HF can provide important information in selecting the most appropriate therapy beyond standard approaches guided by EF phenotypic characterization, especially when specific targeted treatment strategies are indicated,²⁵ provided the diagnostic and/or specific treatment strategies are cost-effective, with favorable benefit risk ratios and are in-line with patient goals. For example, a patient with cardiac amyloidosis requires different treatment strategies than standard HF therapies. The diagnosis of such a patient solely as HFpEF or HFrEF without further work-up to confirm the diagnosis of cardiac amyloidosis may deprive the patient potentially life-saving therapies for amyloidosis.

In clinical practice, the etiology of HF has often been placed into two categories: ischemic and nonischemic cardiomyopathy. However, further diagnostic work up for etiology should be carried out beyond the first step of defining ischemic or nonischemic etiology, especially for dilated, infiltrative, hypertrophic and idiopathic cardiomyopathies.²⁵ Many attempts have been made for morphofunctional classifications of cardiomyopathies in the past.^{25, 26, 28, 102} In this statement, we do not provide recommendations for classifications of specific cardiomyopathies, as we feel those remain outside the scope of this document.

11. Perspective for the Non-cardiologist

The majority of the HF care is provided by non-cardiologists, including general practitioners, internal medicine or family medicine clinicians, hospitalists, emergency room providers, and other specialists. We believe the universal definition will be useful to these clinicians for timely diagnosis and management of patients with HF. Important points for the non-cardiologists are: It is critical to optimally identify and treat patients at-risk for HF to prevent or delay the development of HF; recognize that pre-HF patients, such as asymptomatic patients with elevated natriuretic peptide levels likely will require referral to a cardiologist for further diagnostic and treatment strategies to prevent progression of HF,^{68, 103} that the diagnosis and timely treatment of HF should not be missed or delayed in patients with symptoms and signs of HF, and elevated natriuretic peptide levels or patients with evidence of systemic or pulmonary congestion/ elevated filling pressures, and patients with advanced HF would be considered for referral to HF specialists according to their goals.

Reference List

- (1) Braunwald E. Heart failure. *JACC Heart Fail* 2013 February;1(1):1-20.
- (2) Wagner S, Cohn K. Heart failure. A proposed definition and classification. *Arch Intern Med* 1977 May;137(5):675-8.
- (3) Yancy CW, Jessup M, Bozkurt B et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013 October 15;128(16):e240-e327.
- (4) Ponikowski P, Voors AA, Anker SD et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC
1. *Eur Heart J* 2016 July 14;37(27):2129-200.
- (5) Tsutsui H, Isobe M, Ito H et al. JCS 2017/JHFS 2017 Guideline on Diagnosis and Treatment of Acute and Chronic Heart Failure- Digest Version. *Circ J* 2019 September 25;83(10):2084-184.
- (6) Tan LB, Williams SG, Tan DK, Cohen-Solal A. So many definitions of heart failure: are they all universally valid? A critical appraisal. *Expert Rev Cardiovasc Ther* 2010 February;8(2):217-28.
- (7) Harris P. The problem of defining heart failure. *Cardiovasc Drugs Ther* 1994 June;8(3):447-52.
- (8) Denolin H, Kuhn H, Krayenbuehl HP, Loogen F, Reale A. The definition of heart failure. *Eur Heart J* 1983 July;4(7):445-8.
- (9) Cleland JGF, Pellicori P, Clark AL. Prevention or Procrastination for Heart Failure?: Why We Need a Universal Definition of Heart Failure. *J Am Coll Cardiol* 2019 May 21;73(19):2398-400.
- (10) Virani SS, Alonso A, Benjamin EJ et al. Heart Disease and Stroke Statistics-2020 Update: A Report From the American Heart Association. *Circulation* 2020 March 3;141(9):e139-e596.
- (11) Heidenreich PA, Trogdon JG, Khavjou OA et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation* 2011 March 1;123(8):933-44.
- (12) Pinney SP. Disparities in heart failure care: now is the time to focus on health care delivery. *J Am Coll Cardiol* 2014 August 26;64(8):808-10.
- (13) Bozkurt B. Reasons for Lack of Improvement in Treatment With Evidence-Based Therapies in Heart Failure. *J Am Coll Cardiol* 2019 May 21;73(19):2384-7.
- (14) Greene SJ, Fonarow GC, DeVore AD et al. Titration of Medical Therapy for Heart Failure With Reduced Ejection Fraction. *J Am Coll Cardiol* 2019 May 21;73(19):2365-83.

- (15) Bozkurt B, Ezekowitz J. Substance and Substrate: LVEF and Sex Subgroup Analyses of PARAGON-HF and PARADIGM-HF Trials. *Circulation* 2020 February 4;141(5):362-6.
- (16) Vaduganathan M, Claggett BL, Jhund PS et al. Estimating lifetime benefits of comprehensive disease-modifying pharmacological therapies in patients with heart failure with reduced ejection fraction: a comparative analysis of three randomised controlled trials. *Lancet* 2020 July 11;396(10244):121-8.
- (17) Bonow RO, Ganiats TG, Beam CT et al. ACCF/AHA/AMA-PCPI 2011 performance measures for adults with heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Performance Measures and the American Medical Association-Physician Consortium for Performance Improvement. *Circulation* 2012 May 15;125(19):2382-401.
- (18) Heidenreich PA, Fonarow GC, Brethett K et al. 2020 ACC/AHA Clinical Performance and Quality Measures for Adults With Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Performance Measures. *Circ Cardiovasc Qual Outcomes* 2020 November;13(11):e000099.
- (19) Bozkurt B, Hershberger RE, Butler J et al. TEMPORARY NOTICE: 2021 ACC/AHA Key Data Elements and Definitions for Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Data Standards (Writing Committee to Develop Clinical Data Standards for Heart Failure). *J Am Coll Cardiol* 2020 November 26.
- (20) Fiuzat M, Lowy N, Stockbridge N et al. Endpoints in Heart Failure Drug Development: History and Future. *JACC Heart Fail* 2020 June;8(6):429-40.
- (21) Hicks KA, Mahaffey KW, Mehran R et al. 2017 Cardiovascular and Stroke Endpoint Definitions for Clinical Trials. *Circulation* 2018 February 27;137(9):961-72.
- (22) Thygesen K, Alpert JS, Jaffe AS et al. Fourth Universal Definition of Myocardial Infarction (2018). *Circulation* 2018 November 13;138(20):e618-e651.
- (23) McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971 December 23;285(26):1441-6.
- (24) Adamo L, Nassif ME, Novak E, LaRue SJ, Mann DL. Prevalence of lactic acidemia in patients with advanced heart failure and depressed cardiac output. *Eur J Heart Fail* 2017 August;19(8):1027-33.
- (25) Bozkurt B, Colvin M, Cook J et al. Current Diagnostic and Treatment Strategies for Specific Dilated Cardiomyopathies: A Scientific Statement From the American Heart Association. *Circulation* 2016 December 6;134(23):e579-e646.
- (26) Maron BJ, Towbin JA, Thiene G et al. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary

Working Groups; and Council on Epidemiology and Prevention. *Circulation* 2006 April 11;113(14):1807-16.

- (27) Katz AM. Cardiomyopathy of overload. A major determinant of prognosis in congestive heart failure. *N Engl J Med* 1990 January 11;322(2):100-10.
- (28) Arbustini E, Narula N, Tavazzi L et al. The MOGE(S) classification of cardiomyopathy for clinicians. *J Am Coll Cardiol* 2014 July 22;64(3):304-18.
- (29) Yancy CW, Jessup M, Bozkurt B et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation* 2017 August 8;136(6):e137-e161.
- (30) Inker LA, Astor BC, Fox CH et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD. *Am J Kidney Dis* 2014 May;63(5):713-35.
- (31) Blecker S, Katz SD, Horwitz LI et al. Comparison of Approaches for Heart Failure Case Identification From Electronic Health Record Data. *JAMA Cardiol* 2016 December 1;1(9):1014-20.
- (32) Sabbah HN. Silent disease progression in clinically stable heart failure. *Eur J Heart Fail* 2017 April;19(4):469-78.
- (33) Seid MA, Abdela OA, Zeleke EG. Adherence to self-care recommendations and associated factors among adult heart failure patients. From the patients' point of view. *PLoS One* 2019;14(2):e0211768.
- (34) Gislason GH, Rasmussen JN, Abildstrom SZ et al. Persistent use of evidence-based pharmacotherapy in heart failure is associated with improved outcomes. *Circulation* 2007 August 14;116(7):737-44.
- (35) Santos GC, Liljeroos M, Dwyer AA et al. Symptom perception in heart failure: a scoping review on definition, factors and instruments. *Eur J Cardiovasc Nurs* 2020 February;19(2):100-17.
- (36) Zakeri R, Cowie MR. Heart failure with preserved ejection fraction: controversies, challenges and future directions. *Heart* 2018 March;104(5):377-84.
- (37) Felker GM, Thompson RE, Hare JM et al. Underlying causes and long-term survival in patients with initially unexplained cardiomyopathy. *N Engl J Med* 2000 April 13;342(15):1077-84.
- (38) Atherton JJ, Sindone A, De Pasquale CG et al. National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand: Guidelines for the Prevention, Detection, and Management of Heart Failure in Australia 2018. *Heart Lung Circ* 2018 October;27(10):1123-208.
- (39) Abraham WT, Potoska MA, Fiuzat M. Standardized Definitions for Heart Failure: Scientific Expert Panel from the Heart Failure Collaboratory and Academic Research Consortium (HF-ARC). *JACC Heart Fail* 2020.

- (40) Bohm M, Bewarder Y, Kindermann I. Ejection fraction in heart failure revisited- where does the evidence start? *Eur Heart J* 2020 July 1;41(25):2363-5.
- (41) Solomon SD, Claggett B, Lewis EF et al. Influence of ejection fraction on outcomes and efficacy of spironolactone in patients with heart failure with preserved ejection fraction. *Eur Heart J* 2016 February 1;37(5):455-62.
- (42) Solomon SD, Vaduganathan M, Claggett L et al. Sacubitril/Valsartan Across the Spectrum of Ejection Fraction in Heart Failure. *Circulation* 2020 February 4;141(5):352-61.
- (43) Lund LH, Claggett B, Liu J et al. Heart failure with mid-range ejection fraction in CHARM: characteristics, outcomes and effect of candesartan across the entire ejection fraction spectrum. *Eur J Heart Fail* 2018 August;20(8):1230-9.
- (44) Cleland JGF, Bunting KV, Flather MD et al. Beta-blockers for heart failure with reduced, mid-range, and preserved ejection fraction: an individual patient-level analysis of double-blind randomized trials. *Eur Heart J* 2018 January 1;39(1):26-35.
- (45) Zheng SL, Chan FT, Nabeebaccus AA et al. Drug treatment effects on outcomes in heart failure with preserved ejection fraction: a systematic review and meta-analysis. *Heart* 2018 March;104(5):407-15.
- (46) Tsuji K, Sakata Y, Nochioka K et al. Characterization of heart failure patients with mid-range left ventricular ejection fraction-a report from the CHART-2 Study. *Eur J Heart Fail* 2017 October;19(10):1258-69.
- (47) Chioncel O, Lainscak M, Seferovic PM et al. Epidemiology and one-year outcomes in patients with chronic heart failure and preserved, mid-range and reduced ejection fraction: an analysis of the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail* 2017 December;19(12):1574-85.
- (48) van Veldhuisen DJ, Linssen GC, Jaarsma T et al. B-type natriuretic peptide and prognosis in heart failure patients with preserved and reduced ejection fraction. *J Am Coll Cardiol* 2013 April 9;61(14):1498-506.
- (49) Lam CSP, Gamble GD, Ling LH et al. Mortality associated with heart failure with preserved vs. reduced ejection fraction in a prospective international multi-ethnic cohort study. *Eur Heart J* 2018 May 21;39(20):1770-80.
- (50) Rastogi A, Novak E, Platts AE, Mann DL. Epidemiology, pathophysiology and clinical outcomes for heart failure patients with a mid-range ejection fraction. *Eur J Heart Fail* 2017 December;19(12):1597-605.
- (51) Vedin O, Lam CSP, Koh AS et al. Significance of Ischemic Heart Disease in Patients With Heart Failure and Preserved, Midrange, and Reduced Ejection Fraction: A Nationwide Cohort Study. *Circ Heart Fail* 2017 June;10(6).
- (52) Linde C, Curtis AB, Fonarow GC et al. Cardiac resynchronization therapy in chronic heart failure with moderately reduced left ventricular ejection fraction: Lessons from the Multicenter

InSync Randomized Clinical Evaluation MIRACLE EF study. *Int J Cardiol* 2016 January 1;202:349-55.

- (53) Pellikka PA, She L, Holly TA et al. Variability in Ejection Fraction Measured By Echocardiography, Gated Single-Photon Emission Computed Tomography, and Cardiac Magnetic Resonance in Patients With Coronary Artery Disease and Left Ventricular Dysfunction. *JAMA Netw Open* 2018 August 3;1(4):e181456.
- (54) Hunt SA, Baker DW, Chin MH et al. ACC/AHA Guidelines for the Evaluation and Management of Chronic Heart Failure in the Adult: Executive Summary A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1995 Guidelines for the Evaluation and Management of Heart Failure): Developed in Collaboration With the International Society for Heart and Lung Transplantation; Endorsed by the Heart Failure Society of America. *Circulation* 2001 December 11;104(24):2996-3007.
- (55) Baran DA, Grines CL, Bailey S et al. SCAI clinical expert consensus statement on the classification of cardiogenic shock: This document was endorsed by the American College of Cardiology (ACC), the American Heart Association (AHA), the Society of Critical Care Medicine (SCCM), and the Society of Thoracic Surgeons (STS) in April 2019. *Catheter Cardiovasc Interv* 2019 July 1;94(1):29-37.
- (56) Ammar KA, Jacobsen SJ, Mahoney DW et al. Prevalence and prognostic significance of heart failure stages: application of the American College of Cardiology/American Heart Association heart failure staging criteria in the community. *Circulation* 2007 March 27;115(12):1563-70.
- (57) Wagner M, Tiffe T, Morbach C, Gelbrich G, Stork S, Heuschmann PU. Characteristics and Course of Heart Failure Stages A-B and Determinants of Progression - design and rationale of the STAAB cohort study. *Eur J Prev Cardiol* 2017 March;24(5):468-79.
- (58) Morbach C, Gelbrich G, Tiffe T et al. Prevalence and determinants of the precursor stages of heart failure: results from the population-based STAAB cohort study. *Eur J Prev Cardiol* 2020 May 6;2047487320922636.
- (59) Young K, Scott C, Rodeheffer RJ, Chen CH. Preclinical Heart Failure: Evaluation of Long-Term Outcomes in Patients with Stage A and Stage B Heart Failure in the General Population. *Circulation* 134[Supplement 1], A15103-A15193. 3-29-2018.

Ref Type: Abstract

- (60) Correa de Sa DD, Hodge DO, Slusser JP et al. Progression of preclinical diastolic dysfunction to the development of symptoms. *Heart* 2010 April;96(7):528-32.
- (61) Bozkurt B, Aguilar D, Deswal A et al. Contributory Risk and Management of Comorbidities of Hypertension, Obesity, Diabetes Mellitus, Hyperlipidemia, and Metabolic Syndrome in Chronic Heart Failure: A Scientific Statement From the American Heart Association. *Circulation* 2016 December 6;134(23):e535-e578.
- (62) Zinman B, Wanner C, Lachin JM et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med* 2015 November 26;373(22):2117-28.

- (63) Neal B, Perkovic V, Mahaffey KW et al. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med* 2017 August 17;377(7):644-57.
- (64) Wiviott SD, Raz I, Bonaca MP et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2019 January 24;380(4):347-57.
- (65) Packer M, Anker SD, Butler J et al. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med* 2020 August 29.
- (66) McMurray JJV, Solomon SD, Inzucchi SE et al. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med* 2019 November 21;381(21):1995-2008.
- (67) Berg DD, Wiviott SD, Scirica BM et al. Heart Failure Risk Stratification and Efficacy of Sodium-Glucose Cotransporter-2 Inhibitors in Patients With Type 2 Diabetes Mellitus. *Circulation* 2019 November 5;140(19):1569-77.
- (68) Ledwidge M, Gallagher J, Conlon C et al. Natriuretic peptide-based screening and collaborative care for heart failure: the STOP-HF randomized trial. *JAMA* 2013 July 3;310(1):66-74.
- (69) Saunders JT, Nambi V, de Lemos JA et al. Cardiac troponin T measured by a highly sensitive assay predicts coronary heart disease, heart failure, and mortality in the Atherosclerosis Risk in Communities Study. *Circulation* 2011 April 5;123(13):1367-76.
- (70) deFilippi CR, de Lemos JA, Christenson RH et al. Association of serial measures of cardiac troponin T using a sensitive assay with incident heart failure and cardiovascular mortality in older adults. *JAMA* 2010 December 8;304(22):2494-502.
- (71) Avila MS, Ayub-Ferreira SM, de Barros Wanderley MR Jr et al. Carvedilol for Prevention of Chemotherapy-Related Cardiotoxicity: The CECCY Trial. *J Am Coll Cardiol* 2018 May 22;71(20):2281-90.
- (72) Wilcox JE, Fang JC, Margulies KB, Mann DL. Heart Failure With Recovered Left Ventricular Ejection Fraction: JACC Scientific Expert Panel. *J Am Coll Cardiol* 2020 August 11;76(6):719-34.
- (73) Butler J, Braunwald E, Gheorghiade M. Recognizing worsening chronic heart failure as an entity and an end point in clinical trials. *JAMA* 2014 August 27;312(8):789-90.
- (74) Greene SJ, Mentz RJ, Felker GM. Outpatient Worsening Heart Failure as a Target for Therapy: A Review. *JAMA Cardiol* 2018 March 1;3(3):252-9.
- (75) Solomon SD, Claggett B, Packer M et al. Efficacy of Sacubitril/Valsartan Relative to a Prior Decompensation: The PARADIGM-HF Trial. *JACC Heart Fail* 2016 October;4(10):816-22.
- (76) Halliday BP, Wassall R, Lota AS et al. Withdrawal of pharmacological treatment for heart failure in patients with recovered dilated cardiomyopathy (TRED-HF): an open-label, pilot, randomised trial. *Lancet* 2019 January 5;393(10166):61-73.
- (77) Nishimura RA, Otto CM, Bonow RO et al. 2017 AHA/ACC Focused Update of the 2014 AHA/ACC Guideline for the Management of Patients With Valvular Heart Disease: A Report of

the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2017 June 20;135(25):e1159-e1195.

- (78) January CT, Wann LS, Calkins H et al. 2019 AHA/ACC/HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society in Collaboration With the Society of Thoracic Surgeons. *Circulation* 2019 July 9;140(2):e125-e151.
- (79) Pocock SJ, Ariti CA, McMurray JJ et al. Predicting survival in heart failure: a risk score based on 39 372 patients from 30 studies. *Eur Heart J* 2013 May;34(19):1404-13.
- (80) Martin P, DiMartini A, Feng S, Brown R, Jr., Fallon M. Evaluation for liver transplantation in adults: 2013 practice guideline by the American Association for the Study of Liver Diseases and the American Society of Transplantation. *Hepatology* 2014 March;59(3):1144-65.
- (81) Kartoun U, Corey KE, Simon TG et al. The MELD-Plus: A generalizable prediction risk score in cirrhosis. *PLoS One* 2017;12(10):e0186301.
- (82) Vestbo J, Hurd SS, Agusti AG et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2013 February 15;187(4):347-65.
- (83) Nohria A, Lewis E, Stevenson LW. Medical management of advanced heart failure. *JAMA* 2002 February 6;287(5):628-40.
- (84) Simonneau G, Gatzoulis MA, Adatia I et al. Updated clinical classification of pulmonary hypertension. *J Am Coll Cardiol* 2013 December 24;62(25 Suppl):D34-D41.
- (85) Shah SJ, Katz DH, Selvaraj S et al. Phenomapping for novel classification of heart failure with preserved ejection fraction. *Circulation* 2015 January 20;131(3):269-79.
- (86) Stevenson LW. Design of therapy for advanced heart failure. *Eur J Heart Fail* 2005 March 16;7(3):323-31.
- (87) Pitt B, Remme W, Zannad F et al. Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003 April 3;348(14):1309-21.
- (88) Effect of ramipril on mortality and morbidity of survivors of acute myocardial infarction with clinical evidence of heart failure. The Acute Infarction Ramipril Efficacy (AIRE) Study Investigators. *Lancet* 1993 October 2;342(8875):821-8.
- (89) Pfeffer MA, Braunwald E, Moye LA et al. Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction. Results of the survival and ventricular enlargement trial. The SAVE Investigators. *N Engl J Med* 1992 September 3;327(10):669-77.

- (90) Levy D, Larson MG, Vasan RS, Kannel WB, Ho KK. The progression from hypertension to congestive heart failure. *JAMA* 1996 May 22;275(20):1557-62.
- (91) Cheng S, Claggett B, Correia AW et al. Temporal trends in the population attributable risk for cardiovascular disease: the Atherosclerosis Risk in Communities Study. *Circulation* 2014 September 2;130(10):820-8.
- (92) Ibrahim NE, Burnett JC, Jr., Butler J et al. Natriuretic Peptides as Inclusion Criteria in Clinical Trials: A JACC: Heart Failure Position Paper. *JACC Heart Fail* 2020 May;8(5):347-58.
- (93) Greene SJ, Fonarow GC, Butler J. Risk Profiles in Heart Failure: Baseline, Residual, Worsening, and Advanced Heart Failure Risk. *Circ Heart Fail* 2020 June;13(6):e007132.
- (94) Hollenberg SM, Warner SL, Ahmad T et al. 2019 ACC Expert Consensus Decision Pathway on Risk Assessment, Management, and Clinical Trajectory of Patients Hospitalized With Heart Failure: A Report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol* 2019 October 15;74(15):1966-2011.
- (95) Chioncel O, Mebazaa A, Harjola VP et al. Clinical phenotypes and outcome of patients hospitalized for acute heart failure: the ESC Heart Failure Long-Term Registry. *Eur J Heart Fail* 2017 October;19(10):1242-54.
- (96) Gheorghiade M, Zannad F, Sopko G et al. Acute heart failure syndromes: current state and framework for future research. *Circulation* 2005 December 20;112(25):3958-68.
- (97) Mebazaa A, Yilmaz MB, Levy P et al. Recommendations on pre-hospital & early hospital management of acute heart failure: a consensus paper from the Heart Failure Association of the European Society of Cardiology, the European Society of Emergency Medicine and the Society of Academic Emergency Medicine. *Eur J Heart Fail* 2015 June;17(6):544-58.
- (98) Pfeffer MA, Swedberg K, Granger CB et al. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet* 2003 September 6;362(9386):759-66.
- (99) Pitt B, Pfeffer MA, Assmann SF et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014 April 10;370(15):1383-92.
- (100) Solomon SD, McMurray JJV, Anand IS et al. Angiotensin-Neprilysin Inhibition in Heart Failure with Preserved Ejection Fraction. *N Engl J Med* 2019 October 24;381(17):1609-20.
- (101) McMurray JJ, Packer M, Desai AS et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014 September 11;371(11):993-1004.
- (102) Elliott P, Andersson B, Arbustini E et al. Classification of the cardiomyopathies: a position statement from the European Society Of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J* 2008 January;29(2):270-6.

- (103) Mueller C, McDonald K, de Boer RA et al. Heart Failure Association of the European Society of Cardiology practical guidance on the use of natriuretic peptide concentrations. *Eur J Heart Fail* 2019 June;21(6):715-31.
- (104) Armstrong PW, Pieske B, Anstrom KJ et al. Vericiguat in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med* 2020 May 14;382(20):1883-93.
- (105) Teerlink JR, Diaz R, Felker GM et al. Cardiac Myosin Activation with Omecamtiv Mecarbil in Systolic Heart Failure. *N Engl J Med* 2020 November 13.
- (106) Anker SD, Butler J, Filippatos GS et al. Evaluation of the effects of sodium-glucose co-transporter 2 inhibition with empagliflozin on morbidity and mortality in patients with chronic heart failure and a preserved ejection fraction: rationale for and design of the EMPEROR-Preserved Trial. *Eur J Heart Fail* 2019 October;21(10):1279-87.
- (107) Williams DM, Evans M. Dapagliflozin for Heart Failure with Preserved Ejection Fraction: Will the DELIVER Study Deliver? *Diabetes Ther* 2020 October;11(10):2207-19.
- (108) Stevenson LW, Pagani FD, Young JB et al. INTERMACS profiles of advanced heart failure: the current picture. *J Heart Lung Transplant* 2009 June;28(6):535-41.

Figure Legends

Figure 1. Universal Definition of Heart Failure

Figure 2. Stages in the Development and Progression of Heart Failure

Figure 3. New Classification of Heart Failure According to Left Ventricular Ejection Fraction

Tables

Table 1. Heart Failure Definitions in Contemporary Clinical Practice Guidelines

| | |
|-------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ACCF/AHA (2013) ³ | HF is a complex clinical syndrome that results from any structural or functional impairment of ventricular filling or ejection of blood. The cardinal manifestations of HF are dyspnea and fatigue, which may limit exercise tolerance, and fluid retention, which may lead to pulmonary and/or splanchnic congestion and/or peripheral edema. Some patients have exercise intolerance but little evidence of fluid retention, whereas others complain primarily of edema, dyspnea, or fatigue. |
| ESC (2016) ⁴ | HF is a clinical syndrome characterized by typical symptoms (e.g. breathlessness, ankle swelling and fatigue) that may be accompanied by signs (e.g. elevated jugular venous pressure, pulmonary crackles and peripheral edema) caused by a structural and/or functional cardiac abnormality, resulting in a reduced cardiac output and/or elevated intracardiac pressures at rest or during stress. |
| JCS/JHFS (2017) ⁵ | HF is a clinical syndrome consisting of dyspnea, malaise, swelling and/or decreased exercise capacity due to the loss of compensation for cardiac pumping function due to structural and/or functional abnormalities of the heart |

Table 2. Summary of Heart Failure Inclusion Criteria for Recent Clinical Trials-HFrEF

| Trial Name | Age, NYHA Class | LVEF (%) | Natriuretic Peptides | HF Hospitalization or other |
|--------------------------------------|---------------------------------------|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| PARADIGM-HF ¹⁰¹ | Age ≥18 years NYHA II-IV | LVEF <35% | If previous hospitalization, BNP ≥100 pg/ml or NTproBNP ≥400 pg/ml If no previous hospitalization, BNP ≥150 pg/ml or NT-proBNP ≥600 pg/ml | within previous 12 months |
| VICTORIA ¹⁰⁴ | Age ≥18 years NYHA II-IV | LVEF <45% | within past 30 days: NSR, BNP >300 pg/ml, NT-proBNP >1,000 pg/ml AF BNP >500 pg/ml; NT-proBNP >1,600 pg/ml | within 6 months or outpatient IV diuretics for HF within 3 months |
| DAPA-HF ⁶⁶ | Age ≥18 years NYHA II-IV | LVEF ≤40% | If HF hospitalization within 12 months: NT-proBNP ≥400pg/mL If no hospitalization, NT-proBNP ≥600pg/mL AF NT-proBNP ≥900pg/mL | Diagnosis of HF for at least 2 months |
| EMPEROR-Reduced ⁶⁵ | Age ≥18 years NYHA II-IV | LVEF ≤40% | LVEF ≤30%, NT-proBNP ≥600pg/mL (NSR) or ≥1200pg/mL in AF LVEF 31–35%, NT-proBNP ≥1000pg/mL (NSR) or ≥2000pg/mL in AF LVEF 36–40%, NT-proBNP ≥2500pg/mL (NSR) or ≥5000pg/mL in AF LVEF <40% but HF hospitalization within 12months, NT-proBNP ≥600pg/mL (NSR) or ≥1200pg/mL in AF | NYHA Class II-IV at least 3 months |
| GALACTIC-HF ¹⁰⁵ | Age ≥18 and < 85 years, NYHA II-IV | LVEF ≤ 35% | NT-proBNP ≥400pg/mL (NSR) or ≥1200pg/mL in AF; or BNP ≥125 pg/ml (NSR) ≥375 pg/ml | currently hospitalized for HF (inpatients) or had either made an urgent visit to the emergency department or been hospitalized for HF within 12 months (outpatients) |

ACEI= Angiotensin converting enzyme inhibitor, AF= Atrial fibrillation, ARB= Angiotensin receptor blocker, ARNI= Angiotensin receptor-neprilysin inhibitor, CV= Cardiovascular, HF=heart failure, LVEF= Left Ventricular ejection fraction, MRA= Mineralocorticoid receptor antagonist, NYHA= New York Heart Association, NSR= Normal sinus rhythm,

Table. 3 Summary of Heart Failure Inclusion Criteria for Recent Clinical Trials-HFpEF

| Trial Name | Age, NYHA Class | LVEF (%) | Natriuretic Peptides | HF Hospitalization |
|-----------------------------------------|------------------------------------------------------|-----------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| TOPCAT ⁹⁹ | Age≥50 years NYHA II-IV | LVEF≥45% | BNP ≥100 pg/ml or NT-proBNP ≥360 pg/ml | within previous 12 months, w/ management of HF a major component |
| PARAGON-HF ¹⁰⁰ | Age≥50 years NYHA II-IV | LVEF≥45% and LAE LVH | If NSR, NT-proBNP >200 pg/ml If AF: >600 pg/ml Or If Np previous hospitalization and If NSR: NT-proBNP >300 pg/ml, if AF: NT-proBNP >900 pg/ml | within previous 9 months |
| EMPEROR-Preserved ¹⁰⁶ | Age≥18 years NYHA II-IV (at least 3 months) | LVEF >40% (no prior history of LVEF≤40%) | NT-proBNP >300pg/mL in NSR or >900pg/mL in AF | within 12 months OR evidence of structural changes (LAE or increased LVM) on echo |
| DELIVER ¹⁰⁷ | Age≥40 years NYHA II-IV | (LVEF >40% and evidence of structural heart disease (i.e. LAE or LVH) | Elevated natriuretic peptides | medical history HF ≥6 weeks before enrolment with at least intermittent need for diuretic treatment |

AF= Atrial fibrillation, CV= Cardiovascular, ECG= electrocardiogram, Echo=echocardiogram, HF=heart failure, LAE=left atrial enlargement, LVEF= Left Ventricular ejection fraction, LBBB=Left bundle branch block, LVH= left ventricular hypertrophy, LVM=Left ventricular mass, NYHA= New York Heart Association, NSR= Normal sinus rhythm

| Table 4. Selected Classification Frameworks Currently Used for Heart Failure | |
|-------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| NYHA Class ³ | I, II, III, IV based on symptoms severity |
| Ejection fraction ⁴ | HFrEF, HFmrEF, HFpEF based on left ventricular ejection fraction |
| Etiology ²⁵ | Specific etiology of heart failure, e.g. ischemic/nonischemic, valvular, hypertensive, infiltrative cardiomyopathy such as cardiac amyloidosis, peripartum cardiomyopathy, viral myocarditis chemotherapy induced cardiomyopathy |
| Disease Progression (ACCF/AHA) ^{3, 54} | Stages A, B, C, D according to presence of heart failure symptoms and signs and cardiac structural changes |
| MOGES ²⁸ | Morpho-functional phenotype (M), organ(s) involvement (O), genetic inheritance pattern (G), etiological annotation (E) including genetic defect or underlying disease/substrate, and the functional status (S) |
| INTERMACS Profiles for Advanced HF ¹⁰⁸ | Profiles 1 through 7 according to symptoms, functional capacity, hemodynamic stability for patients who are considered for advanced heart failure therapies |

INTERMACS = Interagency Registry for Mechanically Assisted Circulatory Support ¹⁰⁸

MOGE(S) nosology system ²⁸

NYHA= New York Heart Association

Table 5. Current Heart Failure Classifications According to Left Ventricular Ejection Fraction in Contemporary Clinical Practice Guidelines

| Society Name | HF Classification According to LVEF | LVEF | Additional requirements |
|-------------------------------------|--------------------------------------------------------------------------------------------------------------------------|-------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| ACCF/AHA (2013) ³ | • Heart Failure with Reduced Ejection Fraction (HFrEF) | ≤40% | Symptoms and signs |
| | • Heart Failure with Preserved Ejection Fraction (HFpEF) | ≥50% | Symptoms and signs |
| | a) HFpEF, Borderline | 41% to 49% | Symptoms and signs |
| | B) HFpEF, Improved | >40% | Symptoms and signs |
| ESC (2016) ⁴ | • Heart Failure with Reduced Ejection Fraction (HFrEF) | <40 % | Symptoms and signs |
| | • Heart Failure with Mid-Range Ejection Fraction (HFmrEF) | 40-49% | Symptoms and signs, elevated levels of natriuretic peptides and at least one additional criterion of relevant structural heart disease (LVH or LAE) or diastolic dysfunction |
| | • Heart Failure with Preserved Ejection Fraction (HFpEF) | ≥50% | Symptoms and signs, elevated levels of natriuretic peptides and at least one additional criterion of relevant structural heart disease (LVH or LAE) or diastolic dysfunction |
| JCS/JHFS (2017) ⁵ | • Heart Failure with Reduced Ejection Fraction (HFrEF) | <40% | |
| | • Heart Failure with Mid-Range Ejection Fraction (HFmrEF) | 40 to < 50% | |
| | • Heart Failure with Preserved Ejection Fraction (HFpEF) | ≥50% | |
| | • Heart Failure with Preserved Ejection Fraction, improved (HFpEF improved) or Heart failure with recovered EF (HFrecEF) | ≥40 % | |

ACC= American College of Cardiology, AHA= American Heart Association, ESC= European Society of Cardiology, JCS= Japanese Cardiology Society, JHFS= Japanese Heart Failure Society, HF=heart failure, LAE=left atrial enlargement, LVEF= Left Ventricular ejection fraction, LVH= left ventricular hypertrophy

Table 6. Symptoms and Signs of Heart Failure

| Symptoms of Heart Failure | |
|----------------------------------|-------------------------------------------------------------------------------------------|
| Typical | Breathlessness |
| | Orthopnea* |
| | Paroxysmal nocturnal dyspnea* |
| | Reduced exercise tolerance * |
| | Fatigue, tiredness [€] |
| | Ankle swelling* |
| | Inability to exercise * |
| | Swelling of parts of the body other than ankles |
| | Bendopnea |
| Less Typical | Nocturnal cough |
| | Wheezing |
| | Bloated feeling [¥] |
| | Postprandial satiety [¥] |
| | Loss of appetite |
| | Decline in cognitive function, confusion (especially in the elderly) [€] |
| | Depression |
| | Dizziness , syncope [€] |
| Signs of Heart Failure | |
| More Specific | Elevated jugular venous pressure * |
| | Third heart sound * |
| | Summation gallop with third and fourth heart sounds |
| | Cardiomegaly, laterally displaced apical impulse |
| | Hepatojugular reflux |
| | Cheyne Stokes respiration in advanced heart failure [€] |
| Less specific | Peripheral edema (ankle, sacral, scrotal) |
| | Pulmonary rales* |
| | Unintentional weight gain (>2 kg/week) |
| | Weight loss (in advanced HF) with muscle wasting and cachexia |
| | Cardiac murmur |
| | Reduced air entry and dullness to percussion at lung bases suggestive of pleural effusion |
| | Tachycardia, Irregular pulse |
| | Tachypnea |
| | Hepatomegaly / Ascites |
| | Cold extremities [€] |
| | Oliguria |
| | Narrow pulse pressure |

*Commonly used in clinical trials, registries, risk scoring and have been tested for sensitivity and specificity.

[¥] Can be typical in the setting of right heart failure or biventricular failure

[€] Common in low perfusion, low cardiac output states

Table 7. Natriuretic Peptide Levels Supporting Definition of Heart Failure

| | Ambulatory | Hospitalized/ Decompensated |
|-----------------|------------|-----------------------------|
| BNP pg/ml | ≥ 35 | ≥ 100 |
| NT-proBNP pg/ml | ≥ 125 | ≥ 300 |

BNP= Brain natriuretic peptide, NT-proBNP=N-terminal pro-b-type natriuretic peptide

| Causes of elevated natriuretic peptide levels other than primary diagnosis of heart failure | |
|----------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------|
| Cardiovascular Causes | Acute coronary syndrome, myocardial infarction |
| | Pulmonary embolism |
| | Myocarditis |
| | Hypertrophic cardiomyopathy |
| | Valvular heart disease |
| | Congenital heart disease |
| | Atrial or ventricular arrhythmias |
| | Heart contusion, cardiac infiltration or malignancy |
| | Cardioversion, ICD shock |
| | Pericardial disease |
| | Invasive or surgical procedures involving the heart |
| | Pulmonary hypertension, right ventricular failure |
| | Infiltrative cardiomyopathies |
| Noncardiovascular causes | Advanced age |
| | Kidney disease |
| | Critical illnesses including Sepsis syndrome, cytokine syndrome |
| | Ischemic or hemorrhagic stroke |
| | Pulmonary disease (pneumonia, chronic obstructive pulmonary disease) |
| | Liver disease |
| | Severe anemia |
| | Severe metabolic and hormone abnormalities (e.g. thyrotoxicosis, diabetic ketoacidosis, severe burns) |
| Causes of lower natriuretic peptide levels | |
| | Obesity, or increased BMI |
| | Pericardial disease * |

* In certain patients with pericardial disease and effusion natriuretic peptides may be lower and rise after pericardiocentesis.

Figure 1. Universal Definition of Heart Failure




Figure 2. Stages in the Development and Progression of Heart Failure





Figure 3. New Classification of Heart Failure According to Left Ventricular Ejection Fraction



Appendix 1. Author Relationships With Industry and Other Entities

| Writing Committee Member | Employment/ Affiliation | Consultant | Speakers Bureau | Ownership/ Partnership/ Principal | Personal Research | Institutional, Organizational, or Other Financial Benefit | Expert Witness |
|--------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|-----------------------------------|--------------------------|-----------------------------------------------------------|----------------|
| Magdy | Professor of Cardiovascular Medicine Faculty of Medicine, Kasr Al Ainy, Cairo University , Egypt | None | None | None | None | None | None |
| Stamatis Adampoulos | Director, Heart Failure and Transplant Units, Onassis Cardiac Surgery Centre, Athens-Greece | None | None | None | None | None | None |
| Nancy Albert | Associate Chief Nursing Officer, Research and Innovation- Nursing Institute and Clinical Nurse Specialist, Kaufman Center for Heart Failure- Heart, Vascular and Thoracic Institute, Cleveland Clinic, Cleveland OH | Amgen, AstraZeneca, Boston Scientific, Merck, Novartis | None | None | None | None | None |
| Stefan Anker | Department of Cardiology (CVK), Charité - Universitätsmedizin Berlin; Berlin Institute of Health Center for Regenerative Therapies (BCRT); German Centre for Cardiovascular Research (DZHK), Berlin, Germany | Vifor, Bayer, Boehringer Ingelheim, Novartis, Servier, Abbott, Actimed, Cardiac Dimensions, Impulse Dynamics | None | None | Vifor Int and Abbott, | None | None |
| John Atherton | Director of Cardiology, Royal Brisbane and Women's Hospital, Faculty of Medicine, University of Queensland, Brisbane, QLD, Australia 4006 | Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Eli Lilly, Impedimed, Novartis, Otsuka, Vifor | Amgen, AstraZeneca, Boehringer Ingelheim, Eli Lilly, Novartis | None | None | None | None |

| | | Pharma. | | | | | |
|----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|------|------|-----------------------------------------------------------------------------|------|
| Michael Böhm | Klinik für Innere Medizin III, Universitätsklinikum des Saarlandes, Saarland University Kirrberger Str. 1, 66421 Homburg/Saar, Germany | None | Abbott, Amgen, Astra Zeneca, Bayer, Boehringer Ingelheim, Bristol Myers Squibb, Medtronic, Novartis, Servier, Vifor | None | None | Deutsche Forschungsgemei- n- schaft (DFG, SFB-TTR 219, S- 01 | None |
| Biykem Bozkurt | Baylor College of Medicine, Winters Center for Heart Failure Research, Cardiovascular Research Institute and Michael E. DeBakey VA Medical Center Houston, TX USA | Abbott Amgen Baxter Bristol Myers Squibb Liva Nova RELYPSA/Vifo r Pharma Respicardia Registry Steering Committee Sanofi-aventis scPharmaceutic als, Inc | None | None | None | None | None |
| Javed Butler | University of Mississippi Medical Center | Abbott, Adrenomed, Amgen, Applied Therapeutics, Array, Astra Zeneca, Bayer, Boehringer Ingelheim, CVRx, G3 Pharma, Impul se Dynamics, | Novartis, BI- Lilly, Astra Zeneca, Janssen | None | None | None | None |

| | | | | | | | |
|----------------------|------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|------|------|------|------|
| | | Innolife, Janssen, LivaNova, Luitpold, Medtronic, Merck, Novartis, NovoNordisk, Relypsa, Sequana Medical, V- Wave Limited, and Vifor | | | | | |
| Andrew Coats | University of Warwick, UK and Monash University, Australia | Astra Zeneca, Bayer, Boehringer Ingelheim, Menarini, Novartis, Nutricia, Servier, Vifor, Abbott, Actimed, Arena, Cardiac Dimensions, Corvia, CVRx, Enopace, ESN Clever, Faraday, Gore, Impulse Dynamics, Respicardia | None | None | None | None | None |
| Mark Drazner | Clinical Chief of Cardiology, University of Texas Southwestern Medical Center, Department of Internal Medicine/Division of Cardiology | None | None | None | None | None | None |
| G. Michael Felker | Duke University School of Medicine, Durham, NC | Novartis, Amgen, BMS, Cytokinetics, | None | None | None | None | None |

| | | | | | | | |
|----------------------|------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|------|------|----------------|------|
| | | Medtronic, Cardionomic, Innolife, Boehringer-Ingelheim, American Regent, Abbott, Astra-Zeneca, Eidos Therapeutics, Reprieve, and Sequana, and has served on clinical endpoint committees/data safety monitoring boards for Amgen, Merck, Medtronic, EBR Systems, V-Wave, LivaNova, Siemens, and Rocket Pharma | | | | | |
| Gerasimos Filippatos | National and Kapodistrian University of Athens; Department of Cardiology, Athens University Hospital Attikon, Athens, Greece | Medtronic, Vifor, Servier, Novartis, Boehringer Ingelheim, Bayer, Amgen. | None | None | None | European Union | None |
| Mona Fiuzat | Duke University | None | None | None | None | None | None |
| Gregg Fonarow | Division of Cardiology, University of California, Los Angeles, CA, US | Abbott, Amgen, AstraZeneca, Bayer, CHF Solutions, Edwards, | None | None | NIH | None | None |

Universal Definition of Heart Failure

| | | | | | | | |
|-------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|------|-----------------|-------------------------------------------------------------------------------------------------------------------------|------|
| | | Janssen, Medtronic, Merck, Novartis | | | | | |
| Juan Esteban Gomez-Mesa | Head of the Cardiology Service of the Fundación Valle del Lili. Cali, Colombia; Facultad de Medicina of the Universidad Icesi. Cali, Colombia; Coordinator of the Interamerican Council of Heart Failure & Pulmonary Hypertension of the Interamerican Society of Cardiology. | None | None | None | None | None | None |
| Paul Heidenreich | VA Palo Alto Health Care System, Stanford University School of Medicine | None | None | None | None | None | None |
| Teruhiko Imamura | Second Department of Internal Medicine, University of Toyama | None | None | None | None | None | None |
| James Januzzi | Massachusetts General Hospital | Roche Diagnostics, Siemens, Quidel, Abbott, Novartis, Merck, Jana Care, Imbria | None | None | Novartis, Roche | Endpoint committees: AbbVie, Amgen, Bayer, CVRx, Intercept | None |
| Ewa Jankowska | Department of Heart Diseases, Wroclaw Medical University, Poland and Centre for Heart Diseases, University Hospital, Wroclaw, Poland | Bayer, Abbott, Vifor Pharma, Pfizer, Servier, AstraZeneca, Boehringer Ingelheim, Berlin Chemie, Novartis, Cardiac Dimensions | Bayer, Vifor Pharma, Pfizer, Servier, AstraZeneca, Boehringer Ingelheim, Fresenius, Berlin Chemie, Gedeon Richter, Novartis | None | None | Unrestricted research grant for Wroclaw Medical University Co-PI role in the AFFIRM-AHF trial sponsored by Vifor Pharma | None |
| Prateeti Khazanie | The University of Colorado School of Medicine, Division of Cardiology, Department of Medicine, Aurora, Colorado, US | None | None | None | None | None | None |

Universal Definition of Heart Failure

| | | | | | | | |
|-------------------|--------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-------------------------------------------------------|---------------------------------------------------------------------------------------|------|------|
| Koichiro Kinugawa | University of Toyama | None | None | None | None | None | None |
| Carolyn S.P. Lam | National Heart Centre, Singapore & Duke-National University of Singapore | Abbott Diagnostics, Amgen, Applied Therapeutics, AstraZeneca, Bayer, Biofourmis, Boehringer Ingelheim, Boston Scientific, Corvia Medical, Cytokinetics, Darma Inc., Eko.ai Pte Ltd, JanaCare, Janssen Research & Development LLC, Medtronic, Menarini Group, Merck, MyoKardia, Novartis, Novo Nordisk, Radcliffe Group Ltd., Roche Diagnostics, Sanofi, Stealth BioTherapeutics, The Corpus, Vifor Pharma and WebMD Global LLC | None | co-founder & non-executive director of Eko.ai Pte Ltd | Boston Scientific, Bayer, Roche Diagnostics, AstraZeneca, Medtronic, and Vifor Pharma | None | None |

Universal Definition of Heart Failure

| | | | | | | | |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|------|--------------------------------------------|---------------------------------------------------------------------------------|------|
| Yuya Matsue | Department of Cardiovascular Biology and Medicine, Juntendo University Graduate School of Medicine and Cardiovascular Respiratory Sleep Medicine Juntendo University, Graduate School of Medicine | | Otsuka Pharmaceutical Co., Ltd. and Novartis Japan | | Otsuka Pharmaceutical Co. and Pfizer, Inc. | Philips Respironics, ResMed, and Fukuda Denshi | |
| Marco Metra | University of Brescia | Actelion, Amgen, Astra-Zeneca, Bayer, Livanova, Servier, Vifor pharma, WindTree Therapeutics, a s member of Committees for clinical trials or advisory boards | Abbott vascular, Amgen, Edwards Therapeutics, Servier | none | none | None | none |
| Tomohito Ohtani | Department of Cardiovascular Medicine, Osaka University Graduate School of Medicine | None | Bristol-Myers Squibb Company, Daiichi-Sankyo Co.Ltd, Nippon Boehringer Ingelheim Co.Ltd, Otsuka Pharmaceutical Co. Ltd. | None | None | None | None |
| Massimo Francesco Piepoli | Cardiac Unit, Guglielmo da Saliceto Hospital, Piacenza and University of Parma, Italy | Astra Zeneca, Novartis, Servier, Vifor, TRX Italy | None | None | None | None | None |
| Piotr Ponikowski | Department of Heart Diseases, Wroclaw Medical University, Poland; Centre for Heart Diseases, University Hospital, Wroclaw, Poland | Vifor Pharma, Amgen, Servier, Novartis, Bayer, Pfizer, | Vifor Pharma, Amgen, Servier, Novartis, Pfizer, Impulse Dynamics, Renal | None | None | Abbott Vascular – co-PI of the RESHAPE-2 trial Vifor Pharma – PI of the AFFIRM- | None |

Universal Definition of Heart Failure

| | | | | | | | |
|--------------------|-------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| | | Cibiem, Coridea, Impulse Dynamics, Renal Guard Solutions, Boehringer Ingelheim, AstraZeneca, Corvia, BMS, V-wave Limited | Guard Solutions, Boehringer Ingelheim, AstraZeneca, Berlin Chemie, Radcliffe Group; Ono Pharmaceutical | | | AHF trial Vifor Pharma - research grant for the institution Mediacal Research Agency – Poland: grant for IIT | |
| Giuseppe Rosano | IRCCS San Raffaele Roma, Italy | None | None | None | None | None | None |
| Yasushi Sakata | Department of Cardiovascular Medicine, Osaka University Graduate School of Medicine | None | Otsuka Pharmaceutical Co., Ltd. DAIICHI SANKYO COMPANY, LIMITED Novartis Pharma K.K. Nippon Boehringer Ingelheim Co ., Ltd. | None | None | Astellas Pharma Inc. Otsuka Pharmaceutical Co., Ltd. DAIICHI SANKYO COMPANY, LIMITED Medtronic Japan Co.,Ltd. Nippon Boehringer Ingelheim Co ., Ltd. Bristol-Myers Squibb K.K. | None |

Universal Definition of Heart Failure

| | | | | | | | |
|------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|------|------|------|------|
| Petar Seferovic | Vice-president, European Society of Cardiology President, Heart failure Association of the ESC (2018-2020) Academician, Serbian Academy of Sciences and Arts Professor of Cardiology, University of Belgrade Faculty of Medicine and Heart Failure Center, Belgrade University Medical Center President, Heart Failure Society of Serbia | Boehringer Ingelheim, Novartis, Vifor Pharma | Boehringer Ingelheim, Novartis, Medtronic, Abbott, Servier, Astra Zeneca, Respicardia | None | None | None | None |
| Randy C Starling | Employee Cleveland Clinic Heart, Vascular and Thoracic Institute Cleveland Ohio USA Member FDA Circulatory System Devices Panel | Novartis PARAGLIDE steering committee. No honoraria. Cardiac Dimensions Steering Committee Carillon Trial | None | None | None | None | None |
| John R Teerlink | Section of Cardiology, San Francisco Veterans Affairs Medical Center and School of Medicine, University of California San Francisco | Abbott, Amgen, Astra-Zeneca, Bayer, Boehringer-Ingelheim, Bristol-Myers Squibb, Cytokinetics, Daxor, EBR Systems, Lilly, LivaNova, Medtronic, Merck, Novartis, Relypsa, Servier, | None | None | None | None | None |

Universal Definition of Heart Failure

| | | | | | | | |
|----------------------|------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| | | Windtree Therapeutics, ZS Pharma | | | | | |
| Hiroyuki Tsutsui | Department of Cardiovascular Medicine, Faculty of Medical Sciences, Kyushu University | Nippon Boehringer Ingelheim Co., Ltd. Bayer Yakuhin, Ltd. Novartis Pharma K.K. Ono Pharmaceutical Co., Ltd. | MSD K.K. Astellas Pharma Inc. Pfizer Japan Inc. Bristol-Myers Squibb Company Otsuka Pharmaceutical Co., Ltd. Daiichi Sankyo Co., Ltd. Mitsubishi Tanabe Pharma Corporation Nippon Boehringer Ingelheim Co., Ltd. Takeda Pharmaceutical Company Limited Bayer Yakuhin, Ltd. Novartis Pharma K.K. Kowa Pharmaceutical Co. Ltd. Teijin Pharma Ltd. | None | Actelion Pharmaceuticals Japan Ltd. Japan Tobacco Inc. Mitsubishi Tanabe Pharma Corporation Nippon Boehringer Ingelheim Co., Ltd. Daiichi Sankyo Co., Ltd. IQVIA Services Japan Omron Healthcare | Astellas Pharma Inc. Novartis Pharma K.K. Daiichi Sankyo Co., Ltd. Takeda Pharmaceutical Company Limited Mitsubishi Tanabe Pharma Corporation Teijin Pharma Ltd. MSD K.K. | None |
| Orly Vardeny | Minneapolis Veterans Administration Health Care Center and University of Minnesota | Sanofi-Pasteur, Novartis | None | None | NIH, FDA, AstraZeneca, Bayer | American Heart Association | None |
| Kazuhiro Yamamoto | Department of Cardiovascular Medicine, and Endocrinology and | None | Otsuka Pharmaceutical | None | None | Abbott, Otsuka Pharmaceutical | None |

NOTE: This document contains confidential and/or proprietary information, materials or data. It is important to the integrity of the writing process and final work that this information be kept strictly confidential and not disclosed at any time under any circumstance.

Universal Definition of Heart Failure

| | | | | | | | |
|-----------------|-------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------|------|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| | Metabolism Faculty of Medicine Tottori University | | Co. Ltd, Daiichi-Sankyo Co. Ltd., Novartis, | | | Co. Ltd., Medtronic Japan Co., Ltd., Daiichi-Sankyo Co. Ltd., Biotronik Japan Inc., Japan Lifeline Co. Ltd., Mitsubishi Tanabe Pharma Co. Ltd., Fukuda Denshi, Takeda Pharmaceutical Co. Ltd., Novartis, Ono Pharmaceutical Co. Ltd., Boston Scientific Co. Ltd. | |
| Clyde Yancy | Northwestern University, Feinberg School of Medicine | Spousal Employment, Abbott, Inc. | None | None | None | Deputy Editor, JAMA Network | None |
| Shelley Zieroth | Cardiac Sciences Program, University of Manitoba | Abbott, Akcea, Astra Zeneca, Amgen, Alnylam, Boehringer Ingelheim, Eli- Lilly, Merck, Novartis, Otsuka, Pfizer, Servier, Vifor | Amgen, Astra Zeneca, Boehringer Ingelheim, Eli- Lilly, Novartis, Servier, Vifor | None | Novartis | None | None |
| JianZhang | Fuwai Hospital, Chinese Academy of Medical Science and Peking Union Medical College | No | No | No | No | No | No |

Appendix 2. Abbreviations

ACC = American College of Cardiology

ACEI= angiotensin converting enzyme inhibitor

AHA = American Heart Association

ARB= angiotensin receptor blocker

ARNI= Angiotensin receptor-neprilysin inhibitor

BNP= Brain natriuretic peptide

CKD= chronic kidney disease

CV= Cardiovascular

EF= ejection fraction

ESC= European Society of Cardiology

GDMT= Guideline-directed management and therapy

HFA= Heart Failure Association of European Society of Cardiology

HF = heart failure

HFpEF = heart failure with preserved ejection fraction

HFmrEF= heart failure with mid-range ejection fraction

HFrfEF = heart failure with reduced ejection fraction

JHFS= Japanese Heart Failure Association

LV= left ventricular

LVEF= left ventricular ejection fraction

LVF= left ventricular failure

MRA= mineralocorticoid receptor antagonist

Universal Definition of Heart Failure

NT-proBNP= The N-terminal prohormone of brain natriuretic peptide

NYHA= New York Heart Association

RHF= right heart failure

RV= right ventricular

RVF=right ventricular failure

SGLT2i= sodium-glucose cotransporter-2 inhibitors