


Fertility preservation for medical reasons in girls and women: British fertility society policy and practice guideline

Ephia Yasmin^a, Neerujah Balachandren^a, Melanie C. Davies^a, Georgina L. Jones^b, Sheila Lane^c, Raj Mathur^d, Lisa Webber^a and Richard A. Anderson^e ; on behalf of the British Fertility Society

^aDepartment of Women's Health, University College London Hospitals, London, UK; ^bDepartment of Psychology, School of Social Sciences, Leeds Beckett University, Leeds, UK; ^cDepartment of Paediatric Oncology and Haematology, Children's Oxford University Hospitals NHS Foundation Trust, Oxford, UK; ^dCentral Manchester University Hospitals NHS Foundation Trust and Manchester Academic Health Science Centre, Manchester, UK; ^eMRC Centre for Reproductive Health, Queens Medical Research Institute, University of Edinburgh, Edinburgh, UK

ABSTRACT

Fertility preservation in the female poses several challenges due to the invasive nature of the techniques available to achieve it. The guideline aims to bring together the evidence available for the measures for fertility preservation and their outcome. The guideline addresses fertility preservation for medical reasons and includes both oncological and non-oncological causes. The techniques that the guideline considers are: (i) embryo and oocyte cryopreservation; (ii) ovarian tissue cryopreservation; (iii) GnRH agonist suppression and (iv) ovarian transposition. Although ovarian tissue cryopreservation is still considered experimental, the availability of this technique is gaining momentum as more live births from auto-transplanted tissue are reported. The guideline also highlights use of current treatment modalities for benign and malignant conditions that have a better fertility sparing profile. The guideline recommends a multidisciplinary approach in counselling women and girls about the risk to their fertility and available techniques. The role of psychological support in assisting women and girls with decision-making is highlighted. The guideline also highlights the risks associated with these techniques. Women need to be medically fit to undergo invasive procedures. Fertility preservation techniques are appropriate when treatment has curative intent. Fertility preservation is a subject of on-going research on outcomes of different techniques and at the time of publication, studies are still likely to emerge adding to the available literature.

ARTICLE HISTORY

Received 18 September 2017
Accepted 25 December 2017

KEYWORDS

Fertility preservation;
embryo cryopreservation;
oocyte cryopreservation;
ovarian tissue
cryopreservation; oncology

Introduction

Advances in oncology treatments have led to increased survival rates in cancer patients. Efforts to minimize the long-term morbidity of treatments and improve quality of life among cancer survivors are increasingly a priority. One of the most significant late-effects of cancer treatment is the loss of fertility, which is often rated as the most distressing outcome of therapy by cancer survivors (Bastings et al., 2014; Corney & Swinglehurst, 2014; Mancini, Rey, Préau, Malavolti, & Moatti, 2008; Peate, Meiser, Hickey, & Friedlander, 2009; Ruddy et al., 2014). Given that the average age of first time mothers in the UK is continuing to rise (Office of National Statistics, 2016), it is likely that more women will be nulliparous at the time of diagnosis of cancer, adding greater focus to the need for fertility preservation. Fortunately,

techniques used for fertility preservation are becoming more established as evidence grows concerning their efficacy and safety.

A number of guidelines about fertility preservation exist in the worldwide literature. However, comprehensive guidance encompassing all aspects of fertility preservation in the female, from counselling to outcome of techniques of fertility preservation, would further assist clinicians in this rapidly progressing field. This guidance is aimed primarily at clinical practice in the United Kingdom, where provision of specific services is subject to regulation by the Human Fertilisation and Embryology Authority (HFEA) or the Human Tissue Authority (HTA).

The growing utilization of fertility preservation by women and the recognition of the importance of this in their care by the National Institute of Health and

Clinical Excellence (NICE), highlights the need for a framework of care for women who are at risk of loss of fertility (National Institute for Health and Care Excellence [NICE], 2013). The British Fertility Society first produced a strategy for developing policy and practice in fertility services for survivors of cancer in 2003 (British Fertility Society, 2003). Since then the field of fertility preservation has seen significant progress. In 2007, a Joint Collegiate working party representing relevant professional bodies published 'The effects of cancer treatment on reproductive functions' (Royal College of Physicians, The Royal College of Radiologists, & Royal College of Obstetricians and Gynaecologists, 2007). The Scottish Intercollegiate Guideline Network (SIGN) produced guidance on long-term follow-up of survivors of childhood cancer in 2004 and updated this guidance in 2013 (Scottish Intercollegiate Guideline Network [SIGN], 2013). In 2013, the updated NICE guidance on fertility included recommendations on fertility preservation (NICE, 2013). The guidance recommends that discussion about fertility preservation should occur early, NHS eligibility criteria for infertility should not be used for women seeking fertility preservation, and that no lower age limit should be used.

Therefore, this document aims to provide evidence-based guidance to healthcare professionals involved in offering and providing fertility preservation to girls and women.

Scope of the guideline

This guideline evaluates the evidence for the different techniques of fertility preservation in females. The guideline is aimed at oncologists, haematologists, endocrinologists, gynaecologists and all other healthcare professionals, such as nurses, psychologists, counsellors and general practitioners who play important roles in the management of women with cancer considering fertility preservation.

The content of this guideline reflects the fact that the evidence regarding fertility preservation is more robust in the post-pubertal age group, but includes current options for pre-pubertal girls. In addition, we consider those at risk of premature ovarian insufficiency (POI) from non-cancer related conditions and individuals transitioning from female to male gender. Research recommendations are made to encourage the development of evidence in all relevant cohorts.

Elective ('social') fertility preservation is outside the remit of the guideline, although the technical considerations for oocyte freezing are the same.

Background

There are more than 300,000 new cases of cancer every year in the UK (Cancer Research UK, 2014). The most frequently diagnosed malignancies in women of reproductive age are breast cancer, cervical cancer, lymphoma, leukaemia, sarcomas, brain tumours, melanomas and ovarian cancer (Miller et al., 2016).

Chemotherapy, radiotherapy and surgical treatments to the ovary may lead to reduction or loss of ovarian function with consequent subfertility and POI. Novel targeted therapies are increasingly used by oncologists, but their impact on reproductive function is largely unknown. The degree of damage to gonadal function depends on the type and dose of chemotherapy and radiotherapy, the age of the individual and the initial ovarian reserve (Meirow, Biederman, Anderson, & Wallace, 2010). Apart from oncology treatments, a broad range of medical conditions requiring gonadotoxic therapy may compromise fertility potential: for example stem cell transplantation for benign haematological diseases such as sickle cell anaemia and thalassaemia. Surgical therapies for benign gynaecological conditions, such as ovarian cystectomy, can also compromise the ovarian reserve. Genetic conditions, such as fragile X premutation, mosaicism or monosomy X (Turner syndrome), balanced translocation of the X chromosome and autoimmune conditions may also merit consideration for fertility preservation. Fertility preservation in the transgender population of transitioning females to males is another area of increasing interest.

Although many survivors of childhood cancers will subsequently conceive children naturally, higher rates of infertility are reported after some oncology treatments, particularly in those who received radiotherapy to the abdomen and pelvis or high doses of alkylating agents (Barton et al., 2013; Bramswig, Riepenhausen, & Schellong, 2015). The Childhood Cancer Survivor Study cohort shows that the prevalence of subfertility is higher in survivors compared with their siblings, even when ovarian function is maintained (Barton et al., 2013). Increasingly, more precise data are emerging as to which treatments have significant effects on fertility, and which treatments are associated with a better fertility prognosis (Chow et al., 2016).

Embryo cryopreservation has been established in practice for thirty years. Mature oocyte cryopreservation is an alternative option for fertility preservation and is no longer experimental (Practice Committees of the American Society for Reproductive Medicine and the Society for Assisted Reproductive Technology [ASRM], 2013a). Ovarian tissue cryopreservation (OTC) is less widely available, particularly in the UK, and is

still considered experimental, as is *in vitro* maturation (IVM) of immature oocytes (ASRM, 2013a; ASRM, 2013b).

The likelihood of subsequent utilization of stored oocytes, embryos or ovarian tissue is unknown. Many patients will conceive children naturally, whilst others may not be able to use their stored oocytes or embryos. The longer, more extensive experience with sperm cryopreservation in oncology suggests that future utilization of stored material is likely to be low (Agarwal et al., 2005; Cardozo et al., 2015; Pacey & Eiser, 2011), and due consideration to the management of storage banks is important.

Materials and methods

A search of online databases (MEDLINE, EMBASE, Cochrane Library and Central Register of Controlled Trials) was performed using the keywords: fertility preservation, cancer, oncofertility, oncology, embryo cryopreservation, oocyte cryopreservation, ovarian tissue freezing, ovarian transposition, ovarian suppression, GnRH agonists and gonadotoxicity. Searches were carried out for randomized control trials, systematic reviews, existing guidelines, cohorts, case series and case studies until September 2016. Key publications after this date were also identified. Only English language articles were selected, as resources for translations of non-English articles were not available. Peer-reviewed articles, reviews and guidance from professional interdisciplinary bodies were included. Conference abstracts and unpublished studies were excluded. The literature search and decisions regarding study inclusion were undertaken by all authors. The interpretation of final data and quality assessment was undertaken by the co-ordinating author (EY) and reviewed by the guideline group. We identified 20 published guidelines from professional bodies (Table 1).

Grading of evidence

A recommendation for good practice point (GPP) is made based on the consensus of the guidance development group.

The Royal College of Obstetricians and Gynaecologists grading of evidence was used (Table 2).

Results and recommendations

Limitations of available literature

Randomized control trials were lacking for many of the techniques for fertility preservation. Cohort studies,

small non-randomized trials and case series formed the main body of evidence for most of the techniques.

Pregnancy outcome in cancer survivors

Naturally conceived pregnancies in healthy cancer survivors, who were not exposed to irradiation are probably at no higher risk than in the general population. There may be an increased risk of miscarriage should conception occur within a few months of chemotherapy, but not in the longer term (Signorello et al., 2012; Winther et al., 2012). In particular, there is no evidence of increased risk of congenital anomalies, even after ovarian irradiation or exposure to alkylating agents (Signorello et al., 2012).

Several large population-based studies have examined the outcome of pregnancies in survivors of childhood cancer. A British study found that female survivors exposed to abdominal irradiation had a significantly increased risk of preterm delivery and low birth weight (Reulen et al., 2009). An increased risk of miscarriage was associated with abdominal radiotherapy. Live birth amongst survivors were two-thirds lower than expected. Mueller et al. (2009) demonstrated similar findings of preterm birth and low birth weight from the US Childhood Cancer Survivor Study data. Infants born to childhood cancer survivors were more likely to be preterm and have low birth weight. However, reassuringly, no increased risks of malformations or infant death were found, suggesting no increased germ cell mutation. In a Danish study of 472 survivors of childhood and adolescent cancer, no increase in genetic defects was seen in the offspring despite mutagenic chemotherapy and radiotherapy doses to the gonads (Winther et al., 2012).

Pelvic irradiation (including abdomino-pelvic and spinal irradiation in girls) is associated with increased obstetric risks due to poor uterine function, especially when exposure occurs before menarche (Bath et al., 1999; Larsen et al., 2004; Signorello et al., 2012). These risks include early and late miscarriage, prematurity, low birth weight, stillbirth, neonatal haemorrhage, postpartum haemorrhage, and possibly uterine rupture and placental attachment disorders, such as placenta accreta and percreta (Wo & Viswanathan, 2009).

It is outside the scope of this document to consider fully assessment of fitness for pregnancy after cancer, the impact of pregnancy on the risk of recurrence or the impact of a recurrence on pregnancy. However, these issues do need to be addressed prior to embarking on pregnancy. For example maternal cardiac failure precipitated by pregnancy in cancer survivors has

Table 1. List of currently available guidance.

	Authors	Organization	Title	Journal	Country
1	Loren et al. (2013)	American Society of Clinical Oncology (ASCO)	Fertility preservation for patients with cancer: American society of clinical oncology clinical practice guideline update	Journal of Clinical Oncology	United States
2.	Practice Committee of the ASRM (2013a)	American Society for Reproductive Medicine (ASRM)	Fertility preservation in patients undergoing gonadotoxic therapy or gonadectomy: a committee opinion	Fertility and Sterility	United States
3	Practice Committee of the ASRM (2013b)	American Society for Reproductive Medicine (ASRM)	Mature oocyte cryopreservation: A guideline	Fertility and Sterility	United States
4	Practice Committee of the ASRM (2014)	American Society for Reproductive Medicine (ASRM)	Ovarian tissue cryopreservation: A committee opinion	Fertility and Sterility	United States
5	Martinez et al. (2013)	Barcelona-meeting	Cancer and fertility preservation: Barcelona consensus meeting	Gynaecological Endocrinology	Spain
6	Follows et al. (2014)	British Committee for Standards in Haematology	Guidelines for the first line management of classical Hodgkin lymphoma	British Journal of Haematology	United Kingdom
7	Roberts et al. (2015)	Canadian Fertility and Andrology Society (CFAS)	Fertility preservation in reproductive age woman facing gonadotoxic treatments	Canadian Fertility and Andrology Society	Canada
9.	Joshi, et al. (2014)	Center for International Blood and Marrow Transplant Research: Late Effects Working Committee	Clinical guide to fertility preservation in hematopoietic cell transplant recipients	Bone Marrow Transplant	International
10	Cancer Council Australia (2014)	Clinical Oncology Society of Australia [COSA] (2014)	Fertility preservation for AYAs diagnosed with cancer. Guidance for health professionals	https://wiki.cancer.org.au/australia/COSA:AYA_cancer_fertility_preservation	Australia
12	Cardoza et al. (2012)	European Society of Breast Cancer Specialists (EUSOMA)	The European Society of Breast Cancer Specialists recommendations for the management of young women with breast cancer	European Journal of Cancer	Europe
13	van Wolff et al. (2011)	FertiPROTEKT	Fertility preservation in women- a practical guide to preservation techniques and therapeutic strategies in breast cancer,	Archives Gynaecology Obstetrics	Austria, Germany
14	Jadoul & Kim (2012)	International Society of Fertility Preservation (ISFP)	Fertility preservation in young women with haematological malignancies	Journal of Assisted Reproduction and Genetics	International
15	Kim et al. (2012)	ISFP	Recommendations for fertility preservation in patients with lymphoma, leukaemia and breast cancer	Journal of Assisted Reproduction and Genetics	International
16	Klemp & Kim (2012)	ISFP	Fertility preservation in young women with breast cancer	Journal of Assisted Reproduction and Genetics	International
17	Schmidt & Anderson (2012)	ISFP	Recommendations for fertility preservation in patients with Lymphoma	Journal of Assisted Reproduction and Genetics	International
18	National child cancer network [NCCN] (2014)	National child cancer network (NCCN)	Fertility preservation for people with Cancer: A New Zealand Guideline	https://www.starship.org.nz/media/261297/fertility-preservation-for-people-with-cancer-dec-2017.pdf	New Zealand
19	NICE (2013)	National Institute for Health and Care Excellence	Fertility assessment and treatment for people with fertility problems. Clinical guideline 156	https://www.nice.org.uk/guidance/cg156	UK
20	SIGN (2013)	Scottish Intercollegiate Guidelines Network	Long-term follow up of survivors of childhood cancer	http://www.sign.ac.uk/assets/sign132.pdf	UK

been described associated with exposure to anthracyclines, high dose cyclophosphamide or cardiac irradiation, which may include the scatter from local radiotherapy for breast cancer and not just mediastinal

irradiation (Appel, Nielsen, Zerahn, Jensen, & Skagen, 2007; Hudson, 2010). Pregnancy has not been shown to increase the risk of cancer recurrence, including breast cancer, but with the exception of trophoblastic

Table 2. The Royal College of Obstetricians and Gynaecologists grading of reference.

Classification of evidence levels	Grades of recommendations
1++ High-quality meta-analyses, systematic reviews of randomized controlled trials or randomized controlled trials with a very low risk of bias	A. At least one meta-analysis, systematic reviews or randomized controlled trial rated as 1++, and directly applicable to the target population; or a systematic review of randomized controlled trials or a body of evidence consisting principally of studies rated as 1+, directly applicable to the target population and demonstrating overall consistency of results
1+ Well-conducted meta-analyses, systematic reviews of randomized controlled trials or randomized controlled trials with a low risk of bias	
1– Meta-analyses, systematic reviews of randomized controlled trials or randomized controlled trials with a high risk of bias	
2++ High-quality systematic reviews of case-control or cohort studies or high-quality case-control or cohort studies with a very low risk of confounding, bias or chance and a high probability that the relationship is causal	B. A body of evidence including studies rated as 2++ directly applicable to the target population, and demonstrating overall consistency of results; or extrapolated evidence from studies rated as 1++ or 1+
2+ Well-conducted case-control or cohort studies with a low risk of confounding, bias or chance and a moderate probability that the relationship is causal	
2– Case-control or cohort studies with a high risk of confounding, bias or chance and a significant risk that the relationship is not causal	C. A body of evidence including studies rated as 2+ directly applicable to the target population, and demonstrating overall consistency of results; or extrapolated evidence from studies rated as 2++
3 Non-analytical studies, e.g. case reports and case series	
4 Expert opinion	D. Evidence level 3 or 4; or extrapolated evidence from studies rated as 2+

disease. In the UK, these issues will need to be considered as part of the statutory Welfare of the Child assessment when a licensed fertility treatment is planned (such as transfer of a frozen–thawed embryo or fertilization of frozen-warmed oocytes).

Recommendations

- Women should be informed that risk of congenital anomalies or genetic disease is not increased after cancer treatment. C
- Women who received radiotherapy to a field that included the uterus should be informed of the obstetric risks. C
- Women should be informed that there is no evidence of increased risk of cancer recurrence as a result of pregnancy, with most cancers. C

Evaluation of the risk of loss of fertility

Many studies have used amenorrhoea as a primary outcome of the effect of cancer therapy on fertility. The observed risk of amenorrhoea following various treatments for malignancy is elaborately listed in the American Society of Clinical Oncology guideline (Loren et al., 2013).

Amenorrhoea can occur during or shortly after completion of radiotherapy or chemotherapy (Jacobson, Mertens, Spencer, Manatunga, & Howards, 2015) and may be temporary or permanent. However, few studies have had the long-term follow-up necessary to diagnose POI with accuracy. As per the NICE (2013) guidance recommendation, the risks of infertility and treatment options should be discussed with all patients prior to the start of treatment.

Chemotherapy

The risk of infertility is related to the type of chemotherapeutic agent, dose and drug regimen and age at time of treatment. Depletion of the pool of primordial follicles and compromised vascularity of the gonads are thought to be among the gonadotoxic effects of chemotherapy (Morgan, Anderson, Gourley, Wallace, & Spears, 2012). The size of the pre-treatment ovarian reserve will also impact on the degree of ovarian toxicity (Anderson & Cameron, 2011). The gonadotoxicity of many chemotherapeutic agents, especially newer ones, has not been fully evaluated, and their use in combination also complicates assessment of risk. There do not seem to be deleterious long-term effects on the genetic competency of surviving oocytes (Signorello et al., 2012; Winther et al., 2012) and future pregnancies (Signorello et al., 2006).

Recommendation

- The risk of infertility diminished ovarian reserve and POI should be assessed based on age, type and dose of chemotherapy. C

Radiotherapy

The impact of irradiation on the reproductive organs depends on site, dose and age of the patient, fractionation of treatment and adjuvant therapy. Abdominopelvic radiation of 2Gy can cause loss of more than 50% of the primordial pool and radiation doses of 24Gy will usually cause ovarian failure (Anderson et al., 2015). Cranial irradiation causes impairment of hypothalamic–pituitary–gonadal function; radiation

Table 3. Treatment doses of irradiation in common cancers.

Type of cancer	Dose
Hodgkin's lymphoma	20–40 Gy
Non-Hodgkin's lymphoma	30 Gy
NK/T cell lymphoma	50 Gy
Soft tissue sarcoma	50–66 Gy
Skin cancers	18–60 Gy

doses exceeding 50 Gy will usually cause hypogonadism (Gleeson & Shalat, 2005). Patients who undergo bone marrow transplantation have extremely high rates of ovarian failure, ranging from 72 to 100% after total body irradiation (De Bruin et al., 2008). The dose of radiotherapy depends on the diagnosis and stage of the disease. The typical range of doses of radiotherapy in different conditions is listed in Table 3.

Brachytherapy has significant local effects. Vaginal irradiation leads to dryness and vaginal stenosis. Radiation to the uterus can cause vascular, endometrial and myometrial damage leading to tissue fibrosis and restricted uterine capacity and blood flow. The European Society for Human Reproduction and Embryology (ESHRE) guideline on POI highlights abnormal uterine function and the risk of early and late pregnancy complications in women who have received radiotherapy to the uterus (including total body irradiation) (ESHRE, 2015). The effect on uterine function is dose-dependent and also related to age at the time of exposure. Exposure to radiotherapy before puberty causes significant impairment of the development of the uterus (Bath et al., 1999; Larsen et al., 2004) and reduced responsiveness to exogenous sex steroids (Critchley et al., 1992). Implications for pregnancy outcome are discussed in the section above on pregnancy outcome in cancer survivors.

Recommendation

- Patients undergoing pelvic, abdomino-pelvic or cranio-spinal irradiation should be informed of the risk of infertility, depending on the field of direct and scatter exposure. C

Hormone therapy

Women with hormone-sensitive tumours who are advised to take endocrine therapy will have an age-related decline in fertility even if they do not have adjuvant chemotherapy and radiotherapy. The recent Adjuvant Tamoxifen: Longer Against Shorter (ATLAS) study indicated that 10 years of treatment with tamoxifen is superior to a 5 year course, and this will likely result in a larger proportion of breast cancer patients needing to further delay childbearing (Davies et al.,

2013). However, some women may choose to take a break from endocrine therapy in order to have a baby, after discussion with their oncologist.

Recommendation

- The effect of delay in attempting conception due to prolonged endocrine therapy after breast cancer should be borne in mind when advising women about fertility preservation, even if they do not require gonadotoxic therapy. GPP

Techniques of fertility preservation

This section discusses established and as yet experimental approaches to fertility preservation. It should be recognized that the techniques discussed here do not constitute an exhaustive list. In many clinical situations, the patient may have recourse to treatment options that have a lesser impact on fertility than other available treatments. For example this includes medical treatment with selective progesterone receptor modulators and myomectomy rather than hysterectomy for uterine fibroids. For any technique, evidence of safety and efficacy in the context of fertility preservation is needed before its use for this indication.

Cryopreservation of embryos and oocytes

NICE (2013) guidance recommends embryo and oocyte freezing for fertility preservation where a threat exists from oncological treatments or illnesses that compromise fertility.

Cryopreservation of either oocytes or embryos is preceded by controlled ovarian stimulation (COS) and oocyte collection. The possible risks of delay in starting cancer treatment should be included in the discussion of these options with patients, and also the consequences of any complications of the fertility preservation technique. In general, from the start of ovarian stimulation to oocyte retrieval takes approximately 2 weeks, with chemotherapy usually able to start within 48 h of egg retrieval. A longer recovery period may be required before surgery and radiotherapy to the pelvis.

Recommendations

Women in the reproductive age range should be offered fertility preservation if:

- There is a material risk of infertility as a result of the intended treatment. B
- The treatment for the disease has curative intent or there is good prospect for long-term survival. GPP
- The woman is fit for ovarian stimulation and oocyte collection. GPP
- The time required for ovarian stimulation and oocyte collection does not jeopardize prognosis. GPP

Embryo cryopreservation

The most established technique for fertility preservation is cryopreservation of embryos derived from the patient's oocytes and partner or donor sperm. Embryo cryopreservation is suitable for women in a relationship where both partners consent to creation and storage of embryos; counselling should address the requirement for both partners to consent to the use of embryos in any future fertility treatment. In the case of embryos created with donor sperm, future use requires the consent of the woman alone. UK clinics providing licensed fertility treatment are obliged to consider all factors that may affect the welfare of any child conceived through treatment.

Live birth rates from the transfer of thawed cryopreserved embryos are dependent on the age of the woman at the time of oocyte retrieval (Human Fertility and Embryology Authority [HFEA], 2016). It should be noted that these data are from subfertile couples undergoing assisted conception, rather than couples who have embryos cryopreserved for fertility preservation. The live birth rates in women following transfer of pre-chemo/radiotherapy frozen embryos show comparable results when compared with age-matched controls, however, low rates of utilization of stored embryos are also observed (Cardozo et al., 2015). Some of the causes of non-utilization are death and ill-health, natural conceptions and relationship breakdown (Barcroft, Dayoub, & Thong, 2013).

Recommendation

- Women/couples should be advised that embryo cryopreservation is an established technique, with success rates for the transfer of frozen-thawed embryos comparable to those for the transfer of fresh embryos. D

Oocyte cryopreservation

Women who are not in a relationship, or who do not wish to cryopreserve embryos, are offered oocyte

cryopreservation. However, the prospect of relationship breakdown and inability to use embryos in the future must be borne in mind. For this reason, oocyte cryopreservation is an equally important option for women in relationships. The development of vitrification has dramatically changed the success of oocyte cryopreservation, which is no longer experimental (ASRM, 2013a; NICE, 2013). A meta-analysis of the clinical application of oocyte vitrification showed that pregnancy rates did not differ between use of vitrified and fresh oocytes, although the study population was healthy oocyte donors (Cobo, Garcia-Velasco, Domingo, Remohí, & Pellicer, 2013). Indeed, the confidence with oocyte cryopreservation derives mainly from experience with cryopreserved donor oocytes in recipient cycles (Cobo & Diaz, 2011). In one study, oocyte survival after vitrification ranged between 90 and 97%, fertilization rates were between 71 and 79%, clinical pregnancy rates per transfer ranged from 36 to 61% and the clinical pregnancy rate per thawed oocyte was 4.5–12% (Cobo & Diaz, 2011; Cobo et al., 2016). Comparable results have been reported for women whose oocytes were vitrified for social fertility preservation, demonstrating that the outcome of oocyte vitrification depends on the number and quality of oocytes cryopreserved and the age of the woman at cryopreservation (Doyle, Richter, Stillman, Graham, & Tucker, 2016) with an overall vitrified-warmed oocyte to live-born child efficiency of 6.4% (ASRM, 2013a). On-going pregnancy rates in 182 oocyte vitrification/warming cycles were significantly lower in women over 40 years of age compared with younger women (Borini et al., 2006).

Ovarian stimulation and oocyte vitrification can be performed in post-pubertal adolescent girls (Lavery, Islam, Hunt, Carby, & Anderson, 2016) although such interventions in this age group raise specific practical and ethical issues, and it is unknown whether the pregnancy rate obtained from use of cryopreserved oocytes in this age group is the same as for adult women. In the UK, centres performing oocyte retrieval procedures in patients under the age of 18 years require specific inspection by the Care Quality Commission.

A key benefit of oocyte cryopreservation over embryo cryopreservation is that it obviates the need for partner consent to use the stored oocytes in the future.

Recommendation

- Women should be advised that oocyte cryopreservation is an effective technique, which may have a similar success rate to those using fresh oocytes. C

Controlled ovarian stimulation and oocyte retrieval

COS involves administration of exogenous gonadotrophins to promote development and maturation of multiple ovarian follicles, thereby increasing the number of oocytes available for cryopreservation or generation of embryos.

Controlled ovarian stimulation regimens

Safe and effective COS and oocyte retrieval for women undergoing fertility preservation should take into account the following considerations:

- Minimal delay to treatment: the COS regimen should be applicable without delay and aim to elicit a good ovarian response within the shortest possible time, thus minimizing delay in starting cancer therapy or other treatment.
- Ovarian hyperstimulation syndrome (OHSS): OHSS may require hospital assessment and inpatient care, further delaying the start of cancer treatment. Therefore, minimizing the risk of developing OHSS is of particular significance in this group of patients. Oncology patients may also be at increased risk of VTE should OHSS occur. The antagonist protocol with agonist trigger is suitable due to the short stimulation to oocyte collection interval and reduced risk of OHSS.
- Procedural risks: increased risk of haematoma due to thrombocytopenia and infection due to neutropenia, secondary to either the underlying condition or treatments already commenced, should be discussed with the patient and appropriate antibiotic cover provided.
- Anaesthetic risks: patients should be reviewed by an anaesthetist prior to oocyte harvest if any risk factors for anaesthesia are present, e.g. mediastinal mass in lymphoma patients.
- Women with pelvic malignancies (e.g. of the cervix or ovary) may be at risk of dissemination of malignancy and increased risk of bleeding due to vascularity associated with the tumour. It is not known if women with malignancy are at increased risk of VTE during or after COS due to the associated increased oestrogen levels.
- The route of oocyte collection may have to be altered depending on pelvic disease, e.g. abdominal or laparoscopic oocyte collection may be required in women with cervical cancers.

GnRH antagonist regimens exhibit potential advantages over the use of GnRH agonist in fertility

preservation, although there have been no comparative trials. Use of GnRH antagonist enables ovarian stimulation to be started flexibly at any point in the menstrual cycle, thereby minimizing delay in treatment. Retrospective studies indicate that random-start ovarian stimulation protocols are associated with the collection of similar numbers of oocytes, comparable proportions of mature oocytes and similar fertilization rates as conventional early follicular start ovarian stimulation (Cakmak, Katz, Cedars, & Rosen, 2013). There are no studies comparing live birth rates with these regimens. In addition, the risk of OHSS is lower in treatment cycles using GnRH antagonist compared to GnRH agonist. Meta-analysis of 20 trials involving 5141 women undergoing IVF shows a lower rate of moderate or severe OHSS in women receiving GnRH antagonist compared with GnRH agonist (Youssef et al., 2014). A further advantage of GnRH antagonist regimens is that the pituitary gonadotrophs of women receiving GnRH antagonist retain their sensitivity to the initial flare effect of GnRH agonist, allowing the use of GnRH agonist to induce final follicular maturation. This avoids the use of human chorionic gonadotropin (hCG), which has a longer half-life, and hence a greater potential to precipitate OHSS, than endogenous luteinizing hormone. Meta-analysis shows that the use of GnRH agonist 'trigger' for final follicular maturation is associated with a significantly reduced risk of OHSS compared with the use of hCG, both in women receiving fresh embryos and in women donating oocytes (Youssef et al., 2014).

Evidence is lacking on the type of gonadotropin preparation (recombinant FSH vs. purified urinary FSH) and the starting dose of FSH specifically in women requiring fertility preservation. However, evidence from the wider population of women undergoing IVF supports the use of ovarian reserve tests such as serum anti-Mullerian hormone (AMH) or antral follicle count (AFC) to determine the starting dose of FSH (Jayaprakasan et al., 2012; Nelson et al., 2009). There is evidence to suggest that the ovarian response is reduced in women with some cancers (Domingo et al., 2012); hence, the use of a moderately high dose of FSH is justified for ovarian stimulation in these patients.

COS regimens designed specifically to mitigate the effect of elevated oestrogen concentrations during ovarian stimulation have been developed for use in women with breast cancer. This can be extrapolated to other oestrogen-sensitive cancers. Letrozole co-treatment during ovarian stimulation is associated with significantly lower oestradiol concentrations compared

with conventional gonadotropin stimulation, while resulting in similar numbers of oocytes retrieved and cryopreserved (Oktay, Cil, & Bang, 2006). Tamoxifen, a selective oestrogen receptor modulator often used as adjuvant treatment for women with hormone-sensitive breast cancer, has been studied as a means of managing the potential risk posed by elevated oestradiol concentrations during ovarian stimulation in breast cancer. Meirou et al. (2014) reported outcomes of 76 cycles of embryo cryopreservation in women with breast cancer undergoing ovarian stimulation with or without concomitant tamoxifen use. A recent prospective controlled, non-randomized study in women with Stage 3 or less invasive breast cancer showed no increased risk of recurrence in 120 women who underwent fertility preservation using letrozole as part of the ovarian stimulation protocol, over a mean duration of follow-up of 5 years (Kim, Turan, & Oktay, 2016).

A meta-analysis by Friedler, Kok, Gidoni, Raziel, & Ron-El (2012) indicates that the number of retrieved oocytes in women with a cancer diagnosis was significantly lower than in age-matched healthy IVF patients (Friedler et al., 2012). Other authors show conflicting data. Quinn, Cakman, Letourneau, Cedars, & Rosen (2017) demonstrated that response to ovarian stimulation is not impacted by breast cancer. In situations where time permits, consecutive cycles of ovarian stimulation and oocyte retrieval have been reported with the aim of increasing the number of oocytes cryopreserved (Kuang et al., 2014; Turan, Bedoschi, Moy, & Oktay, 2013). This approach may be helpful in women who exhibit a poor response to ovarian stimulation and in whom the initiation of oncological treatment can be delayed without significant adverse effect on prognosis.

Recommendations

- Antagonist protocols should usually be employed as they shorten the duration of treatment and reduce the risk of OHSS. A
- An agonist trigger in an antagonist cycle should be considered as this minimizes the risk of OHSS, unless contraindicated. A
- Consider using an anti-oestrogen [letrozole, clomifene or tamoxifen] during ovarian stimulation in women with oestrogen-sensitive tumours. D

Duration of storage

Data on embryo and sperm storage demonstrate prolonged survival without detriment (Di Santo, Tarozzi, Nadalini, & Borini, 2012). No differences in survival,

fertilization, cleavage, embryo quality, implantation and live-birth rates were observed in slow-frozen oocytes thawed after up to 48 months compared to earlier thaws (Parmegiani et al., 2009). While, it would seem to be appropriate for the duration of storage to be based on the age of the women at the time of treatment, relevant legislation must be taken into account. In situations where there is actual or anticipated 'premature infertility', the Human Fertilisation and Embryology Authority (<https://www.hfea.gov.uk/code-of-practice/17>) allows storage of gametes or embryos for a maximum period of 55 years. Initial storage is for a maximum period of 10 years, and can be extended by further 10 year periods (up to the maximum of 55 years) if it is shown at any time within each extended storage period that the criterion of premature infertility continues to be met. The relevant HFEA consent form is required to be completed by the patient at each extension, along with a medical practitioner's certificate.

When assisted conception with use of the stored oocytes/embryos is planned, the welfare of the child assessment will need to be made according to HFEA guidance. Patients need to be made aware that subsequent NHS-funding for use of oocytes/embryos may depend on meeting criteria for fertility treatment at the time.

Recommendations

- Women/couples should be advised of the length of time their oocytes/embryos can be stored and that this limit is statutory. GPP
- Women/couples should be advised that there is no evidence to indicate that the duration of storage influences the success rate of using of thawed oocytes/embryos. D
- Welfare of the child assessment should be made at the time of use of oocytes/embryos for assisted conception. GPP

Pregnancy outcome using cryopreserved embryos and oocytes

No particular risk has been identified for children born following frozen-thawed embryo replacement (Williams et al., 2013). In a large systematic review of pregnancy outcome after replacement of frozen-thawed embryos and oocytes, the outcome using slow frozen embryos was comparable to that using fresh embryos (Wennerholm et al., 2009) with no increase in congenital abnormalities. Higher birth weights and lower rates of preterm and low birth weights were

observed following use of slow frozen embryos compared with children born after fresh IVF/ICS, and no increased risk of congenital abnormalities was observed in the analysis of the HFEA data carried out by Maheshwari, Raja, & Bhattacharya (2016). However, these large population-based studies were based on the subfertile population and there is no information specific to the use of frozen embryos from cancer patients prior to gonadotoxic treatment.

Recommendations

Women/couples can be advised that:

- Pregnancy outcome using cryopreserved oocytes/embryos reveals no increase in congenital anomalies. D
- Current data do not suggest that pregnancy outcomes from oocyte/embryos from oncology patients differ from other patients. GPP

Ovarian tissue cryopreservation and transplantation

OTC is the only option for pre-pubertal girls (Anderson et al., 2015) and is an option for post-pubertal adolescents and adult women particularly when there is a narrow time window for fertility preservation. It has the disadvantage of being more invasive, requiring laparoscopy, but successful tissue replacement subsequently can restore the endocrine function of the ovary as well as fertility (Jensen et al., 2015). Although OTC is still considered experimental, a number of successful pregnancies have been reported after orthotopic ovarian tissue transplant with a live birth rate of approximately 25% in these series (Donnez et al., 2013; Jensen et al., 2015; Meirow et al., 2016; Van der Ven et al., 2016) approximately half of which were natural conceptions and others after IVF. Live birth has also been reported after heterotopic replacement (Stern et al., 2013) although this is generally regarded as much less successful. Almost all births have resulted from tissue that was slow frozen, but births have also been reported after ovarian tissue vitrification (Donnez & Dolmans, 2015). There is limited experience in application of this technique for restoring fertility when ovarian tissue is cryopreserved in adolescence or childhood (Demeestere et al., 2015).

Primordial follicles are located in the ovarian cortex: therefore, obtaining cortical tissue potentially enables cryopreservation of significant number of oocytes. Ovarian tissue is procured laparoscopically, either by removal of ovarian cortex or unilateral oophorectomy.

Orthotopic transplantation of ovarian tissue involves transplantation of thin, <1.0–1.5 mm thick strips of thawed ovarian tissue into either the medullary portion of the remaining ovary or a nearby peritoneal pocket.

Advantages of this technique include the possibility of natural conception as the ovarian tissue is in close proximity to the Fallopian tube, and importantly the possibility of long-term hormone production avoiding the need for hormone replacement therapy. Disadvantages include the need for general anaesthesia and laparoscopic surgery, and concern about risk of tissue contamination with micrometastatic disease especially in haematological malignancies (Dolmans, Luyckx, Donnez, Andersen, & Greve, 2013; Jensen et al., 2015). To date, there are no case reports of relapse related to re-implanted tissue; however, recurrences may occur many years after original disease irrespective of reproductive tissue re-implantation. It is, therefore, imperative that all cases of disease recurrence following tissue re-implantation are fully investigated and reported; recognizing the difficulty there may be in proving causality.

Resumption of normal ovulatory menstrual cycles has been reported in over 90% of patients within 4–9 months after transplantation, which is consistent with the time necessary to initiate follicular growth and final maturation (Meirow, Roness, Kristensen, & Andersen, 2015). Grafts have been reported to continue functioning for up to 10 years (Donnez & Dolmans, 2015; Jensen et al., 2015) although this is exceptional, with some grafts showing only very short duration of function.

Whilst OTC techniques are available, in adult patients there is no consensus on selection criteria. Most OTC programmes have been carried out on a research basis, with the use of research criteria. The Edinburgh group (Wallace, Smith, Kelsey, Edgar, & Anderson, 2014) published their selection criteria based on age (younger than 35 years), no previous chemotherapy or radiotherapy (mild or non-gonadotoxic chemotherapy acceptable), a realistic chance of surviving for 5 years and a high risk of POI (>50%). To date, these are the only validated criteria but there is a need to update the criteria based on more recent evidence, for instance the availability of accurate measures of ovarian reserve. Where ovarian reserve is reduced, for example with previous chemotherapy, the success from later replacement of ovarian tissue may be diminished. Criteria for offering OTC should logically be aligned with those for oocyte or embryo cryopreservation.

OTC is still not an established treatment and as such, should only be offered by units with relevant clinical and laboratory expertise, protocols and HTA licensing or associated with an established unit using a third-party arrangement. NHS trusts also require local governance requirements to be satisfied before a new procedural technique is introduced.

Recommendations

- OTC can be considered for post-pubertal patients, particularly where there is insufficient time for ovarian stimulation and oocyte cryopreservation. C
- Increasing numbers of live births from transplantation of frozen ovarian tissue suggests that OTC should be considered for pre-pubertal patients. GPP
- These procedures should only be performed in centres with relevant expertise, facilities and HTA licensing. GPP

In vitro maturation

IVM and vitrification of oocytes retrieved from unstimulated ovaries is currently considered an experimental technique that can be offered to those women who cannot delay their cancer treatment. In oncological patients, oocyte maturation rates of 79% and clinical pregnancy rates of 18–30% per embryo transfer have been reported with IVM (Hourvitz et al., 2015). Implantation rates are lower following transfer of embryos derived from *in vitro* matured oocytes compared with fresh oocytes. It may be possible to combine IVM with ovarian tissue cryobanking (Abir et al., 2016).

Recommendation

- IVM should be offered only in specialized units with relevant expertise, facilities and HFEA licensing. GPP

Ovarian suppression

Suppression of ovarian function by administration of GnRH agonists (GnRHa) immediately before and during chemotherapy has been proposed for the preservation of ovarian function. One rationale is that destruction of follicles by chemotherapeutic agents causes a fall in local AMH levels; this, with a rise in FSH, stimulates more follicles to enter the maturation pathway, rendering them more susceptible to the toxic effects of chemotherapy. The rise in FSH is prevented by GnRHa. Suppression of follicle growth and reduced ovarian

blood flow may also contribute, and there is a possibility of direct effects of GnRHa on the ovary.

In recent years, large RCTs have been performed to assess this approach in women with early breast cancer. The most recent meta-analysis, including 12 trials ($n=1231$ women) indicated a significant reduction in the risk of 'premature ovarian failure' (variably defined in the various studies) with OR 0.36, 95% CI 0.23–0.57, although with significant heterogeneity (Lambertini et al., 2015). Eight trials reported rates of amenorrhea at 1 year after chemotherapy, with OR 0.55, 95% CI 0.41–0.73, without significant heterogeneity (Lambertini et al., 2015). None of the studies have assessed the degree of ovarian function 'saved' by GnRHa treatment, for example by use of AMH measurements.

The possible clinical benefits from ovarian protection may be increased fertility, or reduced impact of oestrogen deficiency from delayed POI, although both are likely to also be influenced by post-chemotherapy hormone treatments (tamoxifen, on-going ovarian suppression with GnRHa) that may be indicated in these patients. No studies have been designed to assess these later outcomes, but the meta-analysis indicates that there were more pregnancies in women treated with GnRHa than in those not treated. As this was not a primary outcome for any studies, this result should be treated with caution.

The majority of studies assessing this approach have been in women with breast cancer, although there have been smaller RCTs in women with lymphoma and ovarian cancer which did not show a benefit. These studies were included in an earlier meta-analysis (Del Mastro et al., 2014), which highlighted the lack of power of studies in diseases other than breast cancer. It remains possible that GnRHa treatment might have a beneficial effect in other malignancies (or indeed in other conditions involving cytotoxic therapy) and larger RCTs are required to assess this. The most recent trial to report in Hodgkin lymphoma found no benefit in terms of POI with GnRHa administration although power was low (Demeestere et al., 2016). Whether GnRHa administration during treatment impacts survival has been addressed in three studies in women with breast cancer, and showed no effect (Lambertini et al., 2015). These RCTs have used monthly administration of goserelin or triptorelin. GnRH antagonists have not been used in RCTs.

An additional benefit of GnRHa administration is suppression of menstruation during chemotherapy and the avoidance of heavy menstrual bleeding, which is often due to thrombocytopenia during

chemotherapy. The risk is highest at the nadir of thrombocytopenia. Menstrual suppression can also be achieved with progestagen. GnRHa carries the side-effects of oestrogen-deficiency vasomotor symptoms (hot flushes and night sweats) and a risk of osteoporosis. It would seem inappropriate for women with hormone-sensitive breast cancer to use add-back therapy with low-dose oestrogen patch or oral tibolone to reduce symptom severity. The initial flare effect of GnRHa should be borne in mind when used for menstrual suppression.

Recommendations

- In premenopausal women with early breast cancer consider temporary ovarian suppression with GnRHa started immediately before and continued during chemotherapy as this may partially preserve ovarian function. A
- Women should be advised that it is possible there is a benefit of using GnRHa when other cancers are treated with gonadotoxic chemotherapy. GPP

Ovarian transposition

The ovaries can be protected from radiation injury by moving them out of the radiation field. This technique is called ovarian transposition, ovarian suspension, oophoropexy or ovariopexy and was introduced more than 50 years ago (McCall, Keaty, & Thompson, 1958). The procedure is now performed laparoscopically unless laparotomy is necessary for the primary treatment of the tumour (Al-Asari & Abduljabbar, 2012). Techniques for ovarian transposition vary according to the radiation field shape, size and location, and there are several sites that the ovaries can be moved to (Gershenson, 2005). Instances when this may be performed include cervical, rectal and colon cancer, pelvic Hodgkin lymphoma and Ewing's sarcoma (Gershenson, 2005).

Morice et al. (1998) reported a series of 24 patients who underwent ovarian transposition to the paracolic gutters before radiation for gynaecologic malignancies. The authors concluded that this procedure was a safe and effective method of preserving ovarian function. Complications are rare in the reported literature. The technique involves transposing the ovaries above the pelvic brim and as lateral as possible, aiming to minimize the ovarian dose of radiation (Gubbala et al., 2014). The ovarian ligament is transected and the ovary mobilized. Blood supply is maintained through the infundibulopelvic ligament. Attention should be paid to avoid torsion and extension of the ovarian

vessels which may reduce blood supply to the ovaries, with a subsequent effect on ovarian reserve. The ovaries can be marked with metallic clips for radiologic identification (Al-Badawi et al., 2010).

Ovarian function after ovarian transposition has been measured by gonadotrophins, AMH and fertility outcomes. Retention of ovarian function has been found to be 60–89% after ovarian transposition (Falcone, Attaran, Bedaiwy, & Goldberg, 2014; Gubbala et al., 2014), although as all reports are non-randomized the size of the effect is unclear. The results may also depend on the type of radiotherapy used (Shou, Chen, Chen, Zhu, & Ni, 2015).

Natural pregnancies are possible if tubal function is preserved as part of the ovarian transposition. Morice et al. (1998) reported 16 natural pregnancies after ovarian transposition. In one report of medial ovarian transposition by positioning the ovaries behind the uterus in 11 girls with Hodgkin lymphoma, the authors reported 14 pregnancies over a 15 years follow-up period (Terenziani et al., 2009).

IVF can be carried out following ovarian transposition using abdominal oocyte collection. No difference in fertilization and pregnancy rates have been observed with trans-vaginally and trans-abdominally collected oocytes (Barton, Politch, Benson, Ginsburg, & Garguilo, 2011). Scattered radiation and reduction in ovarian blood supply after transposition appear to be the main factors causing failure of the technique. Adjuvant radiotherapy the extensiveness of primary surgery at the time of ovarian transposition are seen to affect subsequent ovarian function (Gubbala et al., 2014). There is a small risk of pelvic pain due to the potential for ovarian torsion and Fallopian tube infarction (Gubbala et al., 2014).

The potential effect of radiation upon the uterus should be borne in mind when considering ovarian transposition. Uterine damage from doses in the range of 8500 cGy or intra-cavitary brachytherapy essentially precludes successful pregnancy, either naturally or with *in vitro* fertilization (Agorastos, Zafrakas, & Mastrominas, 2009). Where significant uterine damage has occurred, ovarian transposition may allow the patient to retain the potential to have biological children with host surrogacy (Agorastos et al., 2009).

Recommendations

- Consider ovarian transposition to move ovaries away from the field of irradiation. D
- Fallopian tubes should not be transected in order to retain the possibility of natural conception. GPP

Shielding of ovaries during radiotherapy

Aside from surgical approaches to protect the ovaries from irradiation, modification of radiation techniques to achieve the same end have been explored. External shielding is used in an attempt to minimize radiation exposure to ovaries. The entire pelvis cannot be blocked from radiation without the potential for also shielding tumour cells as well. Such blocks have been shown to offer poor ovarian protection owing to the mobility of the pelvic anatomy (Fawcett, Gomez, Barter, Ditchfield, & Set, 2012).

The use of more complex X-ray planning techniques, such as intensity-modulated radiation therapy (IMRT), has also been examined. Although IMRT reduces doses to many organs, its capacity to reduce the dose meaningfully for a tissue as radiosensitive as the oocyte is quite limited, and ovarian exposure during craniospinal irradiation is likely to exceed 3–5 Gy (RBE) even when IMRT is utilized (Pérez-Andújar, Newhauser, Taddei, Mahajan, & Howell, 2013).

Proton therapy is being used for the delivery of craniospinal irradiation in certain centres and has been shown to allow reduction in the radiation dose delivered to visceral organs anterior to the spinal cord (Yoon, 2012); it may be used for targeted pelvic radiotherapy (Fukushima et al., 2015). Proton beam therapy will be available in the UK soon with two centres being set up in the country.

Recommendation

- Ovarian shielding is of limited use to prevent damage from irradiation and generally should not be offered. D

Fertility-sparing surgery

Fertility-sparing surgery is used in gynaecological cancers with the aim of retaining fertility whilst achieving outcomes that are not inferior to more extensive conventional surgery. A well-established fertility-sparing surgical procedure in young women with early-stage cervical cancer is radical trachelectomy. This may be offered in cases of early-stage invasive cervical cancer instead of hysterectomy. A meta-analysis by Willows, Lennox, and Covens (2016) found that among 1238 patients who underwent fertility-sparing surgery for early cervical cancer, 469 pregnancies occurred with a 67% live birth rate. Outcomes after simple trachelectomy or cervical conization were similar. Cervical stenosis and subfertility are common after trachelectomy. The risk of recurrence has not been found to be higher after these procedures (Okugawa et al., 2016).

In women with borderline ovarian tumours (BOT) concern exists about the effect of ovarian stimulation and the risk of microscopic seeding at oocyte collection after ovarian cystectomy. Most pregnancies after conservative treatment of BOT are natural conceptions. Although BOT are associated with good prognosis, caution needs to be exercised because these tumours can relapse and malignant transformation can also occur. Nevertheless, the patient should be informed that stimulation treatment may be associated with an increased risk of relapse. Cryopreservation of ovarian tissue is recommended by certain teams (Fain-Kahn et al., 2009). However, when autologous transplant is carried out at a later date for fertility, re-transplantation of borderline tissue or the de novo development of a borderline tumour cannot be excluded.

In principle, a pregnancy occurring spontaneously after fertility preservation surgery or ovarian stimulation is preferable to cryopreservation of ovarian tissue so that the ovarian reserve is not further reduced by the removal of ovarian tissue. Unfortunately, evidence about fertility preservation in women with BOT is limited. The patient is usually advised to try to conceive spontaneously after fertility-sparing surgery. Ovarian stimulation and oocyte recovery may be considered after ovarian cystectomy if bilateral oophorectomy is planned as a subsequent step and no significant risk has been reported (Daraï et al., 2013). According to a review of the studies, the risk of relapse of a borderline tumour after ovarian stimulation treatment was 19.4% without resulting in mortality (Denschlag et al., 2010). Letrozole during COH may be considered to reduce circulating oestradiol levels. Oocyte retrieval from affected ovaries at the time of oophorectomy can be considered with a view to *in-vitro* maturation of oocytes (Mangili et al., 2016). OTC is also possible although the tissue will only be used if *in vitro* culture of oocytes becomes available.

Recommendations

- Where feasible, fertility-preserving surgery should be considered in selected women with gynaecological malignancies. B
- Fertility preservation for women with BOT may be associated with additional risks and should be discussed with the patient and her oncology team. D

Discussion of oocyte donation, surrogacy, adoption

In discussing the effect of cancer and treatment of fertility, alternatives to oocyte, embryo and OTC should

be explained. These options include oocyte donation and surrogacy, especially where significant radiation damage to the uterus is involved, adoption and life without children.

Recommendation

- Alternatives to fertility preservation should be discussed. GPP

Fertility preservation in special conditions

Women at risk of non-iatrogenic premature ovarian insufficiency

A small number of women are at risk of POI due to chromosomal, genetic or metabolic conditions (such as Turner mosaicism, fragile X syndrome (FMR1) carrier status or galactosaemia).

At present, strategies for fertility preservation in post-pubertal girls who are at risk of POI are limited to OTC alone or in combination with immature oocyte collection from the tissue followed by *in vitro* maturation and vitrification of mature oocytes. In addition, patients with Turner syndrome, FMR1 carrier status and translocation of X chromosome are candidates for preimplantation genetic diagnosis (PGD) (which requires IVF), because of the increased risk of foetal chromosomal or genetic abnormalities.

As the majority of women with Turner syndrome have POI, fertility preservation with oocyte or OTC may only be an option for those who have spontaneous puberty such as Turner mosaicism (Oktay & Bedoschi, 2014). Almost every woman with classic galactosaemia develops POI, although spontaneous conception has been described in this condition (Gubbels et al., 2009; van Erven et al., 2013). Fertility preservation is only likely to be successful in very young pre-pubertal patients by cryopreservation of ovarian tissue.

Benign conditions treated with chemotherapy

Some non-malignant diseases may require treatments that can impact reproductive function. In patients with systemic lupus erythematosus (SLE), the predominant mechanism of infertility is related to medication rather than disease. The incidence of amenorrhoea following cyclophosphamide treatment for SLE ranges from 27 to 60% (Katsifis & Tzioufas, 2004).

A variety of disorders may be treated with bone marrow transplantation or haematopoietic stem cell transplant with preconditioning alkylating chemotherapy with or without radiation, including thalassaemia

major, sickle cell anaemia, aplastic anaemia, Fanconi anaemia and myeloproliferative diseases (Gidoni, Holzer, Tulandi, & Tan, 2008; Tan et al., 2006). The combination of radiation and high-dose chemotherapy required before bone marrow transplant makes recovery of ovarian function in this patient population unlikely, and spontaneous recovery of ovarian function, as defined by a return of menstrual cycles, has been found to occur in only 6% of patients (Griesshammer, Bergmann, & Pearson, 1998). Some of these conditions (e.g. sickle cell disease) will themselves affect ovarian reserve and may reduce the number of oocytes collected, should FP be undertaken. In children and young adults, OTC should also be considered for this group of patients (Lavery et al., 2016).

Pelvic surgery

Surgery for endometriomas, large ovarian dermoid cysts and recurrent benign ovarian cysts is also likely to reduce ovarian reserve. Therefore, women at risk of recurrent ovarian cystectomies should have a discussion about the effect of repeated surgery and the option of oocyte and embryo cryopreservation should be discussed.

Recommendations

- Discuss fertility preservation options with girls and women at risk of premature ovarian insufficiency from non-malignant conditions. GPP
- Advise women and girls that the evidence for the success of fertility preservation for these indications is limited. GPP

Individuals transitioning from female to male gender

The implications for fertility should be discussed with individuals transitioning from female to male gender. The treatment pathway may include administration of GnRHa to suppress the hypothalamic-pituitary ovarian axis in young people and testosterone to induce virilization, with finally oophorectomy with or without hysterectomy. To retain the possibility of becoming a genetic or biological parent, individuals may wish to store gametes, and may decide to postpone sterilizing surgery. Duration of storage will be the same as for other medical indications.

Fertility preservation should be discussed and, if appropriate, carried out as early as possible in this pathway. Oocyte freezing is likely to be the appropriate mode of fertility preservation. The practicalities of

regrafting make it unlikely that OTC would be a viable option in these cases.

There is no report of fertility preservation after testosterone has been given to induce virilization. Although low doses of androgens have been described as an adjunct to ovarian stimulation (Nagels, Rishworth, Siristatidis, & Kroon, 2015), it is not known whether ovarian stimulation and collection of competent oocytes is possible after high doses of androgens prescribed for virilization.

Counselling is strongly recommended in this complex situation; it should also cover the potential use of stored material following transition, i.e. whether the individual would be prepared to carry a pregnancy in the future, or whether surrogacy would be required.

Recommendations

- Discuss fertility preservation with people transitioning from female to male gender. GPP
- If appropriate, fertility preservation should be performed as early as possible in the treatment pathway. GPP
- Offer counselling, both prior to fertility preservation and prior to use of stored material. GPP

Timing of discussion on fertility preservation

It is generally agreed that women with cancer should be given detailed information about the threat that cancer treatment poses to fertility as early as possible in the cancer treatment pathway (De Vos, Smits, & Woodruff, 2014). The optimal time for FP information appears to be shortly after cancer diagnosis and during the cancer treatment planning stage, specifically before the FP consultation appointment (Balthazar, Fritz, & Mersereau, 2011; Garvelink et al., 2015). Information should be provided by both the referring oncologist and fertility team.

Counselling and psychological support in fertility preservation

There is ample evidence that women of reproductive age with cancer have significant concerns about the effect of cancer treatment on their fertility and the means by which they can preserve it (Peate et al., 2012; Tschudin & Bitzer, 2009). The potential to have biological children appears to be high on their list of priorities (Peate et al., 2012). Women find themselves in a situation where they have to come to terms with their diagnosis and make decisions about their cancer and fertility preservation treatments in a very short

period of time. This can be a source of distress and anxiety. However, the provision of guidance for psychological support is difficult because the decision-making process is often complex and hindered by a number of factors with little evidence of psychological interventions that may help support women through the fertility preservation process.

Supporting patients to make decisions about their future fertility depends on a multidisciplinary approach involving clinical, nursing and surgical teams, psychologists and counsellors. Women often express many fears and anxieties, for example over the implications of pursuing fertility preservation alongside cancer diagnosis/cancer recurrence and the potential health of a future child (Wilkes, Coulson, Crosland, Rubin, & Stewart, 2010). A discussion about fertility preservation with a trained counsellor is advocated in existing clinical guidelines (Loren et al., 2013) although many patients do not appear to get access to this type of service (Letourneau, Melisko, Cedars, & Rosen, 2011; Niemasik et al., 2012). Evidence suggests that many women feel that they do not have access to information that would support them with their decision-making and feel 'too rushed' in consultations to initiate or have the fertility discussion (Jones et al., 2016; Jones, et al., 2017; Woodruff, Clayman, & Waimey, 2014). Therefore, it is important that women are given high quality information and the time needed to absorb this information.

Consideration should be paid to the woman's religious or cultural beliefs and individual values. When dealing with adolescent or pre-pubertal girls, identifying these separately from the wishes of family members can be challenging but the patient's autonomy should be preserved where possible (Tschudin et al., 2010).

Many women will experience cancer after having children. The decisional issues for women starting or completing their family are different.

Decision support interventions help patients to think about the choices they face, in ways which will help inform decision-making (Elwyn et al., 2012). Patient decision aids are one such example of a decision support intervention. Their aim is to support people to make decisions between healthcare options. Several patient decision aids exist to support the fertility preservation process for women (Peate et al., 2012). Two are specifically for women of reproductive age with a breast cancer diagnosis (Garvelink et al., 2013; Peate et al., 2012). They have undergone more extensive evaluation and have been found to reduce decisional conflict and regret about the decision made and increase FP knowledge. Recently, a new tool to

support teenage and adult women with any cancer to make fertility preservation decisions has become available, although it is still in the early stages of evaluation (Jones, et al., 2016).

Decisional conflict appears to be lower when fertility preservation is funded compared with when it is not (Mersereau et al., 2013). Surveys of oncologists have found that many express the need for more fertility preservation information to enable them to support women more effectively (Adams, Hill, & Watson, 2013). The implementation of other decision support interventions such as more education and training packages may, therefore, prove beneficial. In particular, those targeted at improving the communication skills needed to improve collaborative decision-making between patient and provider to deliver care that is patient-centred.

The role of psychology and counselling extends beyond decision-making. On-going support needs to be available when patients experience the late effects of cancer treatment and deal with infertility.

Recommendations

- Discuss fertility preservation as early as possible in the cancer treatment pathway. C
- Refer women to a trained counsellor, both prior to fertility preservation and prior to use of stored material. GPP
- Decision-making aids should be provided to support women's fertility preservation decision-making, ideally at cancer diagnosis/treatment planning. GPP

Summary of recommendations

The fertility preservation techniques are summarized in Table 4.

- Women should be informed that risk of congenital anomalies or genetic disease is not increased after cancer treatment. C
- Women who received radiotherapy to a field that included the uterus should be informed of the obstetric risks. C

- Women should be informed that there is no evidence of increased risk of cancer recurrence as a result of pregnancy, with most cancers. C
- The risk of infertility diminished ovarian reserve and premature ovarian insufficiency should be assessed based on age, type and dose of chemotherapy. C
- Patients undergoing pelvic, abdomino-pelvic or cranio-spinal irradiation should be informed of the risk of infertility, depending on the field of direct and scatter exposure. C
- The effect of delay in attempting conception due to prolonged endocrine therapy after breast cancer should be borne in mind when advising women about fertility preservation, even if they do not require gonadotoxic therapy. GPP

Women in the reproductive age range should be offered fertility preservation if:

- There is a material risk of infertility as a result of the intended treatment. B
- The treatment for the disease has curative intent or there is good prospect for long-term survival. GPP
- The woman is fit for ovarian stimulation and oocyte collection. GPP
- The time required for ovarian stimulation and oocyte collection does not jeopardize prognosis. GPP
- Women/couples should be advised that embryo cryopreservation is an established technique, with success rates for the transfer of frozen-thawed embryos comparable to those for the transfer of fresh embryos. D
- Women should be advised that oocyte cryopreservation is an effective technique, which may have a similar success rate to those using fresh oocytes. C
- Antagonist protocols should usually be employed as they shorten the duration of treatment and reduce the risk of OHSS. A
- An agonist trigger in an antagonist cycle should be considered as this minimizes the risk of OHSS, unless contraindicated. A
- Consider using an anti-oestrogen [letrozole, clomifene or tamoxifen] during ovarian stimulation in women with oestrogen-sensitive tumours. D

Table 4. Fertility preservation strategies in females.

Cryopreservation	Surgical	Medical	Delivery of radiation
Embryo cryopreservation	Ovarian transposition	GnRH agonist suppression	Shielding of ovaries
Oocyte cryopreservation	Fertility-sparing surgery		Intensity-modulated radiotherapy
Ovarian tissue cryopreservation (OTC)			Proton beam therapy
OTC with <i>in vitro</i> maturation of oocytes (IVM)			
OTC with <i>in vitro</i> maturation of oocytes (IVM)			

- Women/couples should be advised of the length of time their oocytes/embryos can be stored and that this limit is statutory. GPP
- Women/couples should be advised that there is no evidence to indicate that the duration of storage influences the success rate of using cryopreserved and warmed oocytes/embryos. D
- Welfare of the child assessment should be made at the time of use of oocytes/embryos for assisted conception. GPP

Women/couples can be advised that:

- Pregnancy outcome using cryopreserved oocytes/embryos reveals no increase in congenital anomalies. D
- Current data do not suggest that pregnancy outcomes from oocyte/embryos from oncology patients differ from other patients. GPP
- OTC can be considered for post-pubertal patients, particularly where there is insufficient time for ovarian stimulation and oocyte cryopreservation. C
- Increasing numbers of live births from transplantation of frozen ovarian tissue suggests that OTC should be considered for pre-pubertal patients. GPP
- These procedures should only be performed at centres with relevant expertise, facilities and HTA licensing. GPP
- IVM should be offered only in specialized units with relevant expertise, facilities and HFEA licensing. GPP
- In premenopausal women with early breast cancer consider temporary ovarian suppression with GnRHa started immediately before and continued during chemotherapy as this may partially preserve ovarian function. A
- Women should be advised that it is possible there is a benefit of using GnRHa when other cancers are treated with gonadotoxic chemotherapy. GPP
- Consider ovarian transposition to move ovaries away from the field of irradiation. D
- Fallopian tubes should not be transected during ovarian transposition in order to retain the possibility of natural conception. GPP
- Ovarian shielding is of limited use to prevent damage from irradiation and generally should not be offered. D
- Where feasible, fertility-preserving surgery should be considered in selected women with gynaecological malignancies. B
- Fertility preservation for women with BOT may be associated with additional risks and should be discussed with the patient and her oncology team. D

- Alternatives to fertility preservation should be discussed. GPP
- Discuss fertility preservation options with girls and women at risk of premature ovarian insufficiency from non-malignant conditions. GPP
- Advise women and girls that the evidence for the success of fertility preservation for these indications is limited. GPP
- Discuss fertility preservation with people transitioning from female to male gender. GPP
- If appropriate, fertility preservation should be performed as early as possible in the treatment pathway. GPP
- Offer counselling, both prior to fertility preservation and prior to use of stored material. GPP
- Discuss fertility preservation as early as possible in the cancer treatment pathway. C
- Decision making aids should be provided to support women's fertility preservation decision-making, ideally at cancer diagnosis/treatment planning. GPP

Service organization

Fertility Preservation services should be organized around efficient pathways enabling patients to access the appropriate expertise and treatment in a timely and supportive manner. Local needs and resources should dictate the specific pathways agreed, but close co-operation between relevant disciplines and a patient-centred approach is key. Pathway development requires liaison between the likely sources of referral (such as Oncology and Haematology) and the centres where fertility preservation treatment is carried out. Owing to the time-critical nature of cases, service commissioners should be involved to obtain agreement on funding mechanisms. Patients should be offered follow up in the fertility clinic to advise them of the late effects, provide management and discuss options for fertility.

Embryo and gamete cryopreservation facilities are available in most assisted conception units, which are regulated in the UK by the HFEA. Gonadal tissue cryopreservation requires access to HTA-licensed storage facilities. Any centre providing care to individuals under the age of 18 must meet specific Care Quality Commission or HFEA requirements.

Recommendations for audit

Audit is essential to monitor and improve the quality of care provided to patients who may require fertility

preservation. The paucity of agreed standards is a difficulty that needs to be addressed. There is a role for deriving standards from research. Professional organizations, such as the Fertility Preservation Network, a Special Interest Group within the British Fertility Society, may have a role in developing standards based on surveys of practice and the opinions of clinicians and service users. Useful areas for audit may include the following:

- The proportion of cancer sufferers offered fertility preservation advice in the relevant age group.
- The proportion of cancer sufferers in whom ovarian reserve is measured before and after cancer treatment.
- The uptake of fertility preservation treatment in the relevant age group and among patients offered this option.
- The proportion of patients offered access to evidence-based patient decision aids.
- The time from cancer diagnosis to fertility preservation procedure.

The creation of a central registry in the UK for fertility preservation would be a powerful tool to guide future audit and research. At present, licensed treatments for fertility preservation (embryo and oocyte storage) are regulated by and reported to the HFEA, but the data available are limited. Distinguishing between fertility preservation by indication (cancer diagnosis versus infertility or 'social') would be an important first step to collecting usable data.

Areas for research

This guideline identifies several areas in fertility preservation that require data and information. Some such areas are:

- Prediction of ovarian damage, risk stratification, accurate identification of risk.
- Cohort studies with documentation of pre-treatment parameters (e.g. age and AMH) and treatment given.
- Research in ovarian tissue technologies – risk of re-grafting, micro-metastases and new methods of identifying contaminated tissue.
- Decision-making in fertility preservation.

Conclusions

Challenges in fertility preservation for women with cancer and other conditions include the distress and

vulnerability, the paucity of time, limited quality of evidence for some techniques, and the variation in access to fertility preservation techniques and funding. In some situations, the risk of treatment-related infertility can be difficult to estimate due to the use of new therapies for which there are limited data only. Where the overall risk of infertility is low, patients should be counselled appropriately so that there is no delay to cancer treatment. It is important to emphasise that although some treatments are non-sterilizing, they have the potential to reduce fertility and the reproductive lifespan. Moreover, patients may also require gonadotoxic therapy due to relapse of disease after previous treatment with a regimen with low gonadotoxic therapy. The field of fertility preservation is evolving rapidly and practice will change as the evidence base worldwide develops.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- Abir, R., Ben-Aharon, I., Garor, R., Yaniv, I., Ash, S., Stemmer, S.M., ... Fisch, B. (2016). Cryopreservation of in vitro matured oocytes in addition to ovarian tissue freezing for fertility preservation in paediatric female cancer patients before and after cancer therapy. *Human Reproduction*, *31*, 750–762. doi: [10.1093/humrep/dew007](https://doi.org/10.1093/humrep/dew007).
- Adams, E., Hill, E., & Watson, E. (2013). Fertility preservation in cancer survivors: A national survey of oncologists' current knowledge, practice and attitudes. *British Journal of Cancer*, *108*, 1602–1615. doi: [10.1038/bjc.2013.139](https://doi.org/10.1038/bjc.2013.139).
- Agarwal, A., Ranganathan, P., Kattal, N., Pasqualotto, F., Hallak, J., Khayal, S., & Mascha, E. (2005). Fertility after cancer: A prospective review of assisted reproductive outcome with banked semen specimens. *Fertility and Sterility*, *81*, 342–348. doi: [0.1177/1758834016665078](https://doi.org/0.1177/1758834016665078).
- Agorastos, T., Zafrakas, M., & Mastrominas, M. (2009). Long-term follow-up after cervical cancer treatment and subsequent successful surrogate pregnancy. *Reproductive Biomedicine Online*, *19*, 250–251. doi: [10.1016/S1472-6483\(10\)60080-6](https://doi.org/10.1016/S1472-6483(10)60080-6).
- Al-Asari, S., & Abduljabbar, A. (2012). Laparoscopic ovarian transposition before pelvic radiation in rectal cancer patient: Safety and feasibility. *Annals of Surgical Innovation and Research*, *6*, 9. doi: [10.1186/1750-1164-6-9](https://doi.org/10.1186/1750-1164-6-9).
- Al-Badawi, I.A., Al-Aker, M., AlSubhi, J., Salem, H., Abduljabbar, A., Balaraj, K., & Munkarah, A. (2010). Laparoscopic ovarian transposition before pelvic irradiation: A Saudi tertiary centre experience. *International Journal of Gynaecological Cancer*, *20*, 1082–1086. doi: [10.1111/IGC.0b013e3181e2ace5](https://doi.org/10.1111/IGC.0b013e3181e2ace5).
- Anderson, R.A., & Cameron, D.A. (2011). Pretreatment serum anti-mullerian hormone predicts long term ovarian function and bone mass after chemotherapy for early breast

- cancer. *The Journal of Clinical Endocrinology and Metabolism*, 96, 1336–1343. doi: [10.1210/jc.2010-2582](https://doi.org/10.1210/jc.2010-2582).
- Anderson, R.A., Mitchell, R.T., Kelsey, T.W., Spears, N., Telfer, E.E., & Wallace, W.H. (2015). Cancer treatment and gonadal function: Experimental and established strategies for fertility preservation in children and young adults. *The Lancet Diabetes and Endocrinology*, 3, 556–567. doi: [10.1016/S2213-8587\(15\)00039-X](https://doi.org/10.1016/S2213-8587(15)00039-X).
- Appel, J.M., Nielsen, D., Zerahn, B., Jensen, B.V., & Skagen, K. (2007). Anthracycline-induced chronic cardiotoxicity and heart failure. *Acta Oncologica*, 46, 576–580. doi: [10.1080/02841860601156165](https://doi.org/10.1080/02841860601156165).
- Balthazar, U., Fritz, M., & Mersereau, J. (2011). Fertility preservation: A pilot study to assess pre-visit patient knowledge quantitatively. *Fertility and Sterility*, 95, 1913–1916. doi: [10.1016/j.fertnstert.2011.02.016](https://doi.org/10.1016/j.fertnstert.2011.02.016).
- Barcroft, J., Dayoub, N., & Thong, K.J. (2013). Fifteen year follow-up of embryos cryopreserved in cancer patients for fertility preservation. *Journal of Assisted Reproduction and Genetics*, 30, 1407–1413. doi: [10.1007/s10815-013-0024-z](https://doi.org/10.1007/s10815-013-0024-z).
- Barton, S.E., Najita, J.S., Ginsburgh, E.S., Leisenring, W.M., Stovall, M., Weathers, R.E., ... Diller, L. (2013). Infertility, infertility treatment, and achievement of pregnancy in female survivors of childhood cancer: A report from the Childhood Cancer Survivor Study cohort. *The Lancet Oncology*, 14, 873–881. doi: [10.1016/S1470-2045\(13\)70251-1](https://doi.org/10.1016/S1470-2045(13)70251-1).
- Barton, S.E., Politch, J.A., Benson, C.B., Ginsburg, E.S., & Garguilo, A.R. (2011). Transabdominal follicular aspiration for oocyte retrieval in patients with ovaries inaccessible by transvaginal ultrasound. *Fertility and Sterility*, 95, 1773–1776. doi: [10.1016/j.fertnstert.2011.01.006](https://doi.org/10.1016/j.fertnstert.2011.01.006).
- Bastings, L., Baysal, Ö., Beerendonk, C.C., IntHout, J., Traas, M.A., Verhaak, C.M., ... Nelen, W.L. (2014). Deciding about fertility preservation after specialist counselling. *Human Reproduction*, 29, 1721–1729. doi: [10.1093/humrep/deu136](https://doi.org/10.1093/humrep/deu136).
- Bath, L.E., Critchley, H.O., Chambers, S.E., Anderson, R.A., Kelnar, C.J., & Wallace, W.H. (1999). Ovarian and uterine characteristics after total body irradiation in childhood and adolescence: Response to sex steroid replacement. *British Journal of Obstetrics and Gynaecology*, 106, 1265–1272. doi: [10.1111/j.1471-0528.1999.tb08180.x](https://doi.org/10.1111/j.1471-0528.1999.tb08180.x).
- Borini, A., Lagalla, C., Bonu, M.A., Bianchi, V., Flamigni, C., & Coticchio, G. (2006). Cumulative pregnancy rates resulting from the use of fresh and frozen oocytes: 7 years' experience. *Reproductive Biomedicine Online*, 12, 481–486. doi: [10.1016/S1472-6483\(10\)62002-0](https://doi.org/10.1016/S1472-6483(10)62002-0).
- Bramswig, J.H., Riepenhausen, M., & Schellong, G. (2015). Parenthood in adult female survivors treated for Hodgkin's lymphoma during childhood and adolescence: A prospective, longitudinal study. *The Lancet Oncology*, 16, 667–675. doi: [10.1016/S1470-2045\(15\)70140-3](https://doi.org/10.1016/S1470-2045(15)70140-3).
- British Fertility Society. (2003). A strategy for fertility services for survivors of childhood cancer. *Human Fertility*, 6, 1–40. doi: [10.1080/1464770312331369133](https://doi.org/10.1080/1464770312331369133).
- Cakmak, H., Katz, A., Cedars, M.I., & Rosen, M.P. (2013). Effective method for emergency fertility preservation: Random-start controlled ovarian stimulation. *Fertility and Sterility*, 100, 673–680. doi: [10.1016/j.fertnstert.2013.07.1992](https://doi.org/10.1016/j.fertnstert.2013.07.1992).
- Cancer Research UK. (2014). Cancer Statistics for the UK. Retrieved from <http://www.cancerresearchuk.org/health-professional/cancer-statistics>.
- Cardozo, E.R., Thomson, A.P., Karmon, A.E., Dickinson, K.A., Wright, D.L., & Sabatini, M.E. (2015). Ovarian stimulation and in-vitro fertilization outcomes of cancer patients undergoing fertility preservation compared to age matched controls: A 17-year experience. *Journal of Assisted Reproduction and Genetics*, 32, 587–596. doi: [10.1007/s10815-015-0428-z](https://doi.org/10.1007/s10815-015-0428-z).
- Cardoza, F., Loibl, S., Pagani, O., Graziottin, A., Panizza, P., Martincich, L., ... Harbeck, N. (2012). The European society of breast cancer specialists recommendations for the management of young people with breast cancer. *European Journal of Cancer*, 48, 3355–3377. doi: [10.1016/j.ejca.2012.10.004](https://doi.org/10.1016/j.ejca.2012.10.004).
- Chow, E.J., Stratton, K.L., Leisenring, W.M., Oeffinger, K.C., Sklar, C.A., Donaldson, S.S., ... Green, D.M. (2016). Pregnancy after chemotherapy in male and female survivors of childhood cancer treated between 1970 and 1999: A report from the childhood cancer survivor study cohort. *The Lancet Oncology*, 17, 567–576. doi: [10.1016/S1470-2045\(16\)00086-3](https://doi.org/10.1016/S1470-2045(16)00086-3).
- Clinical Oncology Society of Australia. (2014). Fertility preservation for AYAs diagnosed with cancer: Guidance for health professionals. Retrieved from http://wiki.cancer.org.au/australia/COSA:AYA_cancer_fertility_preservation/Introduction.
- Cobo, A., & Diaz, C. (2011). Clinical application of oocyte vitrification: A systematic review and meta-analysis of randomized controlled trials. *Fertility and Sterility*, 96, 277–285. doi: [10.1016/j.fertnstert.2011.06.030](https://doi.org/10.1016/j.fertnstert.2011.06.030).
- Cobo, A., García-Velasco, J.A., Coello, A., Domingo, J., Pellicer, A., & Remohí, J. (2016). Oocyte vitrification as an efficient option for elective fertility preservation. *Fertility and Sterility*, 105, 755–764. doi: [10.1016/j.fertnstert.2015.11.027](https://doi.org/10.1016/j.fertnstert.2015.11.027).
- Cobo, A., Garcia-Velasco, J.A., Domingo, J., Remohí, J., & Pellicer, A. (2013). Is vitrification of oocytes useful for fertility preservation for age-related fertility decline and in cancer patients? *Fertility and Sterility*, 99, 1485–1495. doi: [10.1016/j.fertnstert.2013.02.050](https://doi.org/10.1016/j.fertnstert.2013.02.050).
- Corney, R.H., & Swinglehurst, A.J. (2014). Young childless women with breast cancer in the UK: A qualitative study of their fertility-related experiences, options, and the information given by health professionals. *Psycho-Oncology*, 23, 20–26. doi: [10.1002/pon.3365](https://doi.org/10.1002/pon.3365).
- Critchley, H.O., Wallace, W.H., Shalet, S.M., Mamtora, H., Higginson, J., & Anderson, D.C. (1992). Abdominal irradiation in childhood; the potential for pregnancy. *British Journal of Obstetrics and Gynaecology*, 99, 392–394. doi: [10.1111/j.1471-0528.1992.tb13755.x](https://doi.org/10.1111/j.1471-0528.1992.tb13755.x).
- Darai, E., Fauvet, R., Uzan, C., Gouy, S., Duvillard, P., & Morice, P. (2013). Fertility and borderline ovarian tumor: A systematic review of conservative management, risk of recurrence and alternative options. *Human Reproduction Update*, 19, 151–166. doi: [10.1093/humupd/dms047](https://doi.org/10.1093/humupd/dms047).
- Davies, C., Pan, H., Godwin, J., Gray, R., Arriagada, R., Raina, V., ... Peto, R. (2013). Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast

- cancer: ATLAS, a randomised trial. *The Lancet*, 381, 805–816. doi: [10.1016/S0140-6736\(12\)61963-1](https://doi.org/10.1016/S0140-6736(12)61963-1).
- De Bruin, M.L., Huisbrink, J., Hauptmann, M., Kuenen, M.A., Ouwens, G.M., Van't Veer, M.B., ... van Leeuwen, F.E. (2008). Treatment-related risk factors for premature menopause following Hodgkin lymphoma. *Blood*, 111, 101–108. doi: [10.1182/blood-2007-05-090225](https://doi.org/10.1182/blood-2007-05-090225).
- Del Mastro, L., Ceppi, M., Poggio, F., Bighin, C., Peccatori, F., Demeestere, I., ... Bruzzi, P. (2014). Gonadotropin-releasing hormone analogues for the prevention of chemotherapy-induced premature ovarian failure in cancer women: Systematic review and meta-analysis of randomized trials. *Cancer Treatment Reviews*, 40, 675–683. doi: [10.1016/j.ctrv.2013.12.001](https://doi.org/10.1016/j.ctrv.2013.12.001).
- Demeestere, I., Brice, P., Peccatori, F.A., Kentos, A., Dupuis, J., Zachee, P., ... De Maertelaer, V. (2016). No evidence for the benefit of gonadotropin-releasing hormone agonist in preserving ovarian function and fertility in lymphoma survivors treated with chemotherapy: Final long-term report of a prospective randomized trial. *Journal of Clinical Oncology*, 22, 2568–2574. doi: [10.1200/JCO.2015.65.8864](https://doi.org/10.1200/JCO.2015.65.8864).
- Demeestere, I., Simon, P., Dedeken, L., Moffa, F., Tsépélidis, S., Brachet, C., ... Ferster, A. (2015). Live birth after autograft of ovarian tissue cryopreserved during childhood. *Human Reproduction*, 30, 2107–2109. doi: [10.1093/humrep/dev128](https://doi.org/10.1093/humrep/dev128).
- Denschlag, D., Wolff, M., Amant, F., Kesic, V., Reed, N., Schneider, A., & Rodolakis, A. (2010). Fertility preservation in borderline ovarian neoplasm (tumor of low malignant potential): Ovarian stimulation and oocyte retrieval after conservative surgery. *Gynecologic and Obstetric Investigation*, 70, 160–165. doi: [10.1159/000316264](https://doi.org/10.1159/000316264).
- De Vos, M., Smits, J., & Woodruff, T.K. (2014). Fertility preservation in women with cancer. *The Lancet*, 384, 1302–1310. doi: [10.1016/S0140-6736\(14\)60834-5](https://doi.org/10.1016/S0140-6736(14)60834-5).
- Di Santo, M., Tarozzi, N., Nadalini, M., & Borini, A. (2012). Human sperm cryopreservation: Update on techniques, effect on DNA integrity, and implications for ART. *Advances in Urology*, 2012, 12. doi: [10.1155/2012/854837](https://doi.org/10.1155/2012/854837).
- Dolmans, M.M., Luyckx, V., Donnez, J., Andersen, C.Y., & Greve, T. (2013). Risk of transferring malignant cells with transplanted frozen-thawed ovarian tissue. *Fertility and Sterility*, 99, 1514–1522. doi: [10.1016/j.fertnstert.2013.03.027](https://doi.org/10.1016/j.fertnstert.2013.03.027).
- Domingo, J., Guillén, V., Ayllón, Y., Martínez, M., Muñoz, E., Pellicer, A., & Garcia-Velasco, J.A. (2012). Ovarian response to controlled ovarian hyperstimulation in cancer patients is diminished even before oncological treatment. *Fertility and Sterility*, 97, 930–934. doi: [10.1016/j.fertnstert.2012.01.093](https://doi.org/10.1016/j.fertnstert.2012.01.093).
- Donnez, J., & Dolmans, M.M. (2015). Ovarian cortex transplantation: 60 reported live births brings the success and worldwide expansion of the technique towards routine clinical practice. *Journal of Assisted Reproduction and Genetics*, 32, 1167–1170. doi: [10.1007/s10815-015-0544-9](https://doi.org/10.1007/s10815-015-0544-9).
- Donnez, J., Dolmans, M.M., Pellicer, A., Diaz-Garcia, C., Sanchez Serrano, M., Schmidt, K.T., ... Andersen, C.Y. (2013). Restoration of ovarian activity and pregnancy after transplantation of cryopreserved ovarian tissue: A review of 60 cases of reimplantation. *Fertility and Sterility*, 99, 1503–1513. doi: [10.1016/j.fertnstert.2013.03.030](https://doi.org/10.1016/j.fertnstert.2013.03.030).
- Doyle, J., Richter, K., Stillman, R., Graham, J., & Tucker, M. (2016). Successful elective and medically indicated oocyte vitrification and warming for autologous in vitro fertilization, with predicted birth probabilities for fertility preservation according to number of cryopreserved oocytes and age at retrieval. *Fertility and Sterility*, 105, 459–466. doi: [10.1016/j.fertnstert.2015.10.026](https://doi.org/10.1016/j.fertnstert.2015.10.026).
- Elwyn, G., Frosch, D., Thomson, R., Joseph-Williams, N., Lloyd, A., Kinnersley, P., ... Barry, M. (2012). Shared decision making: A model for clinical practice. *Journal of General Internal Medicine*, 27, 1361–1367. doi: [10.1007/s11606-012-2077-6](https://doi.org/10.1007/s11606-012-2077-6).
- European Society of Human Reproduction and Embryology. (2015). Guideline on the management of premature ovarian insufficiency. Retrieved from <https://www.eshre.eu/Guidelines-and-Legal/Guidelines/Management-of-premature-ovarian-insufficiency.aspx>.
- Fain-Kahn, V., Poirot, C., Uzan, C., Prades, M., Gouy, S., Genestie, C., ... Morice, P. (2009). Feasibility of ovarian cryopreservation in borderline ovarian tumours. *Human Reproduction*, 24, 850–855. doi: [10.1093/humrep/den459](https://doi.org/10.1093/humrep/den459).
- Falcone, T., Attaran, M., Bedaiwy, M.A., & Goldberg, J.M. (2014). Ovarian function preservation in the cancer patient. *Fertility and Sterility*, 81, 243–257. doi: [10.1016/j.fertnstert.2003.06.031](https://doi.org/10.1016/j.fertnstert.2003.06.031).
- Fawcett, S.L., Gomez, A.C., Barter, S.J., Ditchfield, M., & Set, P. (2012). More harm than good? The anatomy of misguided shielding of the ovaries. *The British Journal of Radiology*, 85, 442–447. doi: [10.1259/bjr/25742247](https://doi.org/10.1259/bjr/25742247).
- Follows, G., Ardeshtna, K.M., Barrington, S.F., Culligan, D.J., Hoskin, P.J., Linch, D., ... Wimperis, J.Z. (2014). Guidelines for the first line management of classical Hodgkin lymphoma. *British Journal of Haematology*, 166, 34–49. doi: [10.1111/bjh.12878](https://doi.org/10.1111/bjh.12878).
- Friedler, S., Kok, O., Gidoni, Y., Raziell, A., & Ron-El, R. (2012). Ovarian response to stimulation for fertility preservation in women with malignant disease: A systematic review and meta-analysis. *Fertility and Sterility*, 97, 125–133. doi: [10.1016/j.fertnstert.2011.10.014](https://doi.org/10.1016/j.fertnstert.2011.10.014).
- Fukushima, H., Fukushima, T., Sakai, A., Suzuki, R., Kobayashi, C., Oshiro, Y., ... Sumazakia, R. (2015). Tailor-made treatment combined with proton beam therapy for children with genitourinary/pelvic rhabdomyosarcoma. *Reports of Practical Oncology and Radiotherapy*, 20, 217–222. doi: [10.1016/j.rpor.2014.12.003](https://doi.org/10.1016/j.rpor.2014.12.003).
- Garvelink, M.M., ter Kuile, M.M., Bakker, R.M., Geense, W.J., Jenninga, E., Louwé, L.A., ... Stiggelbout, A.M. (2015). Women's experiences with information provision and deciding about fertility preservation in the Netherlands: 'Satisfaction in general, but unmet needs'. *Health Expectations*, 18, 956–968. doi: [10.1111/hex.12068](https://doi.org/10.1111/hex.12068).
- Garvelink, M.M., ter Kuile, M.M., Fischer, M.J., Louwé, L.A., Hilders, C.G., Kroep, J.R., & Stiggelbout, A.M. (2013). Development of a decision aid about fertility preservation for women with breast cancer in the Netherlands. *Journal of Psychosomatic Obstetrics and Gynaecology*, 34, 170–188. doi: [10.3109/0167482X.2013.851663](https://doi.org/10.3109/0167482X.2013.851663).
- Gershenson, D.M. (2005). Fertility-sparing surgery for malignancies in women. *Journal of the National Cancer Institute Monographs*, 34, 43–47. doi: [10.1093/jncimonographs/lgi011](https://doi.org/10.1093/jncimonographs/lgi011).

- Gidoni, Y., Holzer, H., Tulandi, T., & Tan, S.L. (2008). Fertility preservation in patients with non-oncological conditions. *Reproductive Biomedicine Online*, 16, 792–800. doi: [10.1016/S1472-6483\(10\)60144-7](https://doi.org/10.1016/S1472-6483(10)60144-7).
- Gleeson, H., & Shalat, S.M. (2005). The impact of cancer therapy on the endocrine system in survivors of childhood brain tumours. *Endocrinology Related Cancer*, 24, 589–602. doi: [10.1677/erc.1.00779](https://doi.org/10.1677/erc.1.00779).
- Griesshammer, M., Bergmann, L., & Pearson, T. (1998). Fertility, pregnancy and the management of myeloproliferative disorders. *Baillieres Clinical Haematology*, 11, 859–874. doi: [10.1016/S0140-6736\(97\)08247-0](https://doi.org/10.1016/S0140-6736(97)08247-0).
- Gubbala, K., Laios, A., Gallos, I., Pathiraja, P., Haldar, K., & Ind, T. (2014). Outcomes of ovarian transposition in gynaecological cancers; a systematic review and meta-analysis. *Journal of Ovarian Research*, 7, 69. doi: [10.1186/1757-2215-7-6](https://doi.org/10.1186/1757-2215-7-6).
- Gubbels, C.S., Kuppens, S.M., Bakker, J.A., Konings, C.J., Wodzig, K.W., Der Velden, M.G., ... Rubio-Gozalbo, M.E. (2009). Pregnancy in classic galactosemia despite undetectable anti-Mullerian hormone. *Fertility and Sterility*, 91, 1213–1296. doi: [10.1016/j.fertnstert.2008.12.031](https://doi.org/10.1016/j.fertnstert.2008.12.031).
- Hourvitz, A., Yerushalmi, G.M., Maman, E., Raanani, H., Elizur, S., Brengauz, M., ... Meirow, D. (2015). Combination of ovarian tissue harvesting and immature oocyte collection for fertility preservation increases preservation yield. *Reproductive Biomedicine Online*, 31, 497–505. doi: [10.1016/j.rbmo.2015.06.025](https://doi.org/10.1016/j.rbmo.2015.06.025).
- Hudson, M.M. (2010). Reproductive outcomes for survivors of childhood cancer. *Obstetrics and Gynecology*, 116, 1171–1183. doi: [10.1097/AOG.0b013e3181f87c4b](https://doi.org/10.1097/AOG.0b013e3181f87c4b).
- Human Fertility and Embryology Authority. (2016). Fertility treatment 2014, trends and figures. Retrieved from http://www.hfea.gov.uk/docs/HFEA_Fertility_treatment_Trends_and_figures_2014.pdf.
- Jacobson, M.H., Mertens, A.C., Spencer, J.B., Manatunga, A.K., & Howards, P.P. (2015). Menses resumption after cancer treatment-induced amenorrhea occurs early or not at all. *Fertility and Sterility*, 105, 765–772. doi: [10.1016/j.fertnstert.2015.11.020](https://doi.org/10.1016/j.fertnstert.2015.11.020).
- Jadoul, P., & Kim, S.S. (2012). Fertility considerations in young women with hematological malignancies. *Journal of Assisted Reproduction and Genetics*, 29, 470–487. doi: [10.1007/s10815-012-9792-0](https://doi.org/10.1007/s10815-012-9792-0).
- Jayaprakasan, K., Chan, Y., Islam, R., Haoula, Z., Hopkisson, J., Coomarasamy, A., & Fenning, R. (2012). Prediction of in vitro fertilization outcome at different antral follicle count thresholds in a prospective cohort of 1,012 women. *Fertility and Sterility*, 98, 657–663. doi: [10.1016/j.fertnstert.2012.05.042](https://doi.org/10.1016/j.fertnstert.2012.05.042).
- Jensen, A.K., Kristensen, S.G., Macklon, K.T., Jeppesen, J.V., Fedder, J., Ernst, E., & Anderson, C.Y. (2015). Outcomes of transplantations of cryopreserved ovarian tissue to 41 women in Denmark. *Human Reproduction*, 30, 2838–2845. doi: [10.1093/humrep/dev230](https://doi.org/10.1093/humrep/dev230).
- Jones, G.L., Hughes, J., Greenfield, D., Lacey, A., Slade, P., Skull, J., ... Ledger, W. (2016). What factors influence the fertility preservation treatment decision-making process in women with cancer? The qualitative findings of the PREFER study. European shared medical decision making conferences. (Abstract). Retrieved from <https://smdm.confex.com/smdm/16BEC/webprogram/Paper9835.html>.
- Jones, G.L., Hughes, J., Mahmoodi, N., Smith, E., Skull, J., & Ledger, W. (2017). What factors hinder the decision-making process for women with cancer and contemplating fertility preservation? *Human Reproduction Update*, 23, 1–25. doi: [10.1093/humupd/dmx009](https://doi.org/10.1093/humupd/dmx009).
- Jones, G.L., Hughes, J., Mahmoodi, N., Greenfield, D., Brauten-Smith, G., Skull, J., ... Bekker, H.L. (2017). Observational study of the development and evaluation of a fertility preservation patient decision aid for teenage and adult women diagnosed with cancer: the Cancer, Fertility and Me research protocol. *BMJ Open*, 7, e013219. doi: [10.1136/bmjopen-2016-013219](https://doi.org/10.1136/bmjopen-2016-013219).
- Joshi, S., Savani, B., Chow, E., Gilleece, M., Halter, J., Jacobsohn, D., ... Majhail, N. (2014). Clinical guide to fertility preservation in hematopoietic cell transplant recipients. *Bone Marrow Transplantation*, 49, 477–484. doi: [10.1038/bmt.2013.211](https://doi.org/10.1038/bmt.2013.211).
- Katsifis, G.E., & Tzioufas, A.G. (2004). Ovarian failure in systemic lupus erythematosus patients treated with pulsed intravenous cyclophosphamide. *Lupus*, 13, 673–678. doi: [10.1191/0961203304lu2012oa](https://doi.org/10.1191/0961203304lu2012oa).
- Kim, J., Turan, V., & Oktay, K. (2016). Long-term safety of letrozole and gonadotropin stimulation for fertility preservation in women with breast cancer. *The Journal of Clinical Endocrinology and Metabolism*, 101, 1364–1371. doi: [10.1210/jc.2015-3878](https://doi.org/10.1210/jc.2015-3878).
- Kim, S.S., Donnez, J., Barri, P., Pellicer, A., Patrizio, P., Rosenwaks, Z., ... Dolmans, M.M. (2012). Recommendations for fertility preservation in patients with lymphoma, leukemia, and breast cancer. *Journal of Assisted Reproduction and Genetics*, 29, 465–468. doi: [10.1007/s10815-012-9786-y](https://doi.org/10.1007/s10815-012-9786-y).
- Klemp, J.R., & Kim, S.S. (2012). Fertility preservation in young women with breast cancer. *Journal of Assisted Reproduction and Genetics*, 29, 469–472. doi: [10.1007/s10815-012-9791-1](https://doi.org/10.1007/s10815-012-9791-1).
- Kuang, Y., Chen, Q., Hong, Q., Lyu, Q., Ai, A., Fu, Y., & Shoham, Z. (2014). Double stimulations during the follicular and luteal phases of poor responders in IVF/ICSI programmes (Shanghai protocol). *Reproductive Biomedicine Online*, 29, 684–691. doi: [10.1016/j.rbmo.2014.08.009](https://doi.org/10.1016/j.rbmo.2014.08.009).
- Lambertini, M., Ceppi, M., Poggio, F., Peccatori, F.A., Jr. Azim, H.A., Ugolini, D., ... Del Mastro, L. (2015). Ovarian suppression using luteinizing hormone-releasing hormone agonists during chemotherapy to preserve ovarian function and fertility of breast cancer patients: A meta-analysis of randomized studies. *Annals of Oncology*, 26, 2408–2419. doi: [10.1093/annonc/mdv374](https://doi.org/10.1093/annonc/mdv374).
- Larsen, E.C., Schmiegelow, K., Rechnittzer, C., Loft, A., Muller, J., & Andersen, A.N. (2004). Radiotherapy at a young age reduces uterine volume of childhood cancer survivors. *Acta Obstetrica Et Gynecologica Scandinavica*, 83, 96–102. doi: [10.1111/j.1600-0412.2004.00332.x](https://doi.org/10.1111/j.1600-0412.2004.00332.x).
- Lavery, S.A., Islam, R., Hunt, J., Carby, A., & Anderson, R.A. (2016). The medical and ethical challenges of fertility preservation in teenage girls: A case series of sickle cell anaemia patients prior to bone marrow transplant. *Human Reproduction*, 31, 1501–1507. doi: [10.1093/humrep/dew084](https://doi.org/10.1093/humrep/dew084).
- Letourneau, J.M., Melisko, M.E., Cedars, M.I., & Rosen, M.P. (2011). A changing perspective: Improving access to

- fertility preservation. *Nature Reviews Clinical Oncology*, 8, 56–60. doi: [10.1038/nrclinonc.2010.133](https://doi.org/10.1038/nrclinonc.2010.133).
- Loren, A.W., Mangu, P.B., Beck, L.N., Brennan, L., Magdalinski, A.J., Partridge, A.H., ... Oktay, K. (2013). Fertility preservation for patients with cancer: American society of clinical oncology clinical practice guideline update. *Journal of Clinical Oncology*, 31, 2500–2510. doi: [10.1200/JCO.2013.49.2678](https://doi.org/10.1200/JCO.2013.49.2678).
- Maheshwari, A., Raja, E.A., & Bhattacharya, S. (2016). Obstetric and perinatal outcomes after either fresh or thawed frozen embryo transfer: An analysis of 112,432 singleton pregnancies recorded in the human fertilisation and embryology authority anonymized dataset. *Fertility and Sterility*, 106, 1703–1708. doi: [10.1016/j.fertnstert.2016.08.047](https://doi.org/10.1016/j.fertnstert.2016.08.047).
- Mancini, J., Rey, D., Préau, M., Malavolti, L., & Moatti, J.P. (2008). Infertility induced by cancer treatment: Inappropriate or no information provided to majority of French survivors of cancer. *Fertility and Sterility*, 90, 1616–1625. doi: [10.1016/j.fertnstert.2007.08.064](https://doi.org/10.1016/j.fertnstert.2007.08.064).
- Mangili, G., Somigliana, E., Giorgione, V., Martinelli, F., Filippi, F., Petrella, M.C., ... Peccatori, F. (2016). Fertility preservation in women with borderline ovarian tumours. *Cancer Treatment Reviews*, 49, 13–24. doi: [10.1016/j.ctrv.2016.06.010](https://doi.org/10.1016/j.ctrv.2016.06.010).
- Martinez, F., Devesa, M., Coroleu, B., Tur, R., González, R., Boada, M., ... Barri, P.N. (2013). Cancer and fertility preservation: Barcelona consensus meeting. *Gynecological Endocrinology*, 29, 285–291. doi: [10.3109/09513590.2012.743019](https://doi.org/10.3109/09513590.2012.743019).
- McCall, M.L., Keaty, E.C., & Thompson, J.D. (1958). Conservation of ovarian tissue in the treatment of carcinoma of the cervix with radical surgery. *American Journal of Obstetrics and Gynecology*, 75, 590–600. doi: [10.1016/0002-9378\(58\)90614-8](https://doi.org/10.1016/0002-9378(58)90614-8).
- Meirow, D., Biederman, H., Anderson, R.A., & Wallace, W.H. (2010). Toxicity of chemotherapy and radiation on female reproduction. *Clinical Obstetrics and Gynecology*, 53, 727–739. doi: [10.1097/GRF.0b013e3181f96b54](https://doi.org/10.1097/GRF.0b013e3181f96b54).
- Meirow, D., Raanani, H., Maman, E., Paluch-Shimon, S., Shapira, M., Cohen, Y., ... Kaufman, B. (2014). Tamoxifen co-administration during controlled ovarian hyperstimulation for in vitro fertilization in breast cancer patients increases the safety of fertility-preservation treatment strategies. *Fertility and Sterility*, 102, 488–495. doi: [10.1016/j.fertnstert.2014.05.017](https://doi.org/10.1016/j.fertnstert.2014.05.017).
- Meirow, D., Ra'anani, H., Shapira, M., Brenghausen, M., Derech Chaim, S., Aviel-Ronen, S., ... Dor, J. (2016). Transplantations of frozen-thawed ovarian tissue demonstrate high reproductive performance and the need to revise restrictive criteria. *Fertility and Sterility*, 106, 467–474. doi: [10.1016/j.fertnstert.2016.04.031](https://doi.org/10.1016/j.fertnstert.2016.04.031).
- Meirow, D., Roness, H., Kristensen, S.G., & Andersen, C.Y. (2015). Optimizing outcomes from ovarian tissue cryopreservation and transplantation; activation versus preservation. *Human Reproduction*, 30, 2453–2456. doi: [10.1093/humrep/dev210](https://doi.org/10.1093/humrep/dev210).
- Mersereau, J.E., Goodman, L.R., Deal, A.M., Gorman, J.R., Whitcomb, B.W., & Su, H.I. (2013). To preserve or not to preserve: How difficult is the decision about fertility preservation? *Cancer*, 119, 4044–4050. doi: [10.1002/cncr.28317](https://doi.org/10.1002/cncr.28317).
- Miller, K.D., Siegel, R.L., Lin, C.C., Mariotto, A.B., Kramer, J.L., Rowland, J.H., ... Jemal, A. (2016). Cancer treatment and survivorship statistics, 2016. *CA Cancer Journal for Clinicians*, 66, 271–289. doi: [10.3322/caac.21349](https://doi.org/10.3322/caac.21349).
- Morgan, S., Anderson, R.A., Gourley, C., Wallace, W.H., & Spears, N. (2012). How do chemotherapeutic agents damage the ovary? *Human Reproduction Update*, 18, 525–535. doi: [10.1093/humupd/dms022](https://doi.org/10.1093/humupd/dms022).
- Morice, P., Thiam-Ba, R., Castaine, D., Haie-Meder, C., Gerbault, A., Pautier, P., ... Michel, G. (1998). Fertility results after ovarian transposition for pelvic malignancies treated by external irradiation or brachytherapy. *Human Reproduction*, 13, 660–666. doi: [10.1093/humrep/13.3.660](https://doi.org/10.1093/humrep/13.3.660).
- Mueller, B.A., Chow, E.J., Kaminen, A., Daling, J.R., Fraser, A., Wiggins, C.L., ... Drews-Botsch, C. (2009). Pregnancy outcomes in female childhood and adolescent cancer survivors: A linked cancer-birth registry analysis. *Archives of Pediatric and Adolescent Medicine*, 163, 879–886. doi: [10.1001/archpediatrics.2009.112](https://doi.org/10.1001/archpediatrics.2009.112).
- Nagels, H.E., Rishworth, J.R., Siristatidis, C.S., & Kroon, B. (2015). Androgens (dehydroepiandrosterone or testosterone) for women undergoing assisted reproduction. *Cochrane Database of Systematic Reviews*, CD009749. doi: [10.1002/14651858.CD009749.pub2](https://doi.org/10.1002/14651858.CD009749.pub2).
- National child cancer network. (2014). Fertility preservation for people with cancer: A New Zealand guideline. Retrieved from https://www.starship.org.nz/media/261297/fertility_preservation_for_people_with_cancer_-_a_new_zealand_guideline_april_2014final.pdf.
- National Institute for Health and Care Excellence. (2013). Fertility assessment and treatment for people with fertility problems. Retrieved from <https://www.nice.org.uk/guidance/cg156>.
- Nelson, S.M., Yates, R.W., Lyall, H., Jamieson, M., Traynor, I., Gaudoin, M., ... Fleming, R. (2009). Anti-Mullerian hormone-based approach to controlled ovarian stimulation for assisted conception. *Human Reproduction*, 24, 867–875. doi: [10.1093/humrep/den480](https://doi.org/10.1093/humrep/den480).
- Niemasik, E.E., Letourneau, J., Dohan, D., Katz, A., Melisko, M., Rugo, H., & Rosen, M. (2012). Patient perceptions of reproductive health counseling at the time of cancer diagnosis: A qualitative study of female California cancer survivors. *Journal of Cancer Survivorship*, 6, 324–332. doi: [10.1007/s11764-012-0227-9](https://doi.org/10.1007/s11764-012-0227-9).
- Office of National Statistics. (2016). Births by parents' characteristics in England and Wales: 2015. Retrieved from <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/livebirths/bulletins/birthsbyparentscharacteristicsinenglandandwales/2015>.
- Oktay, K., & Bedoschi, G. (2014). Oocyte cryopreservation for fertility preservation in postpubertal female children at risk for premature ovarian failure due to accelerated follicle loss in Turner syndrome or cancer treatments. *Journal of Pediatric and Adolescent Gynecology*, 27, 342–346. doi: [10.1016/j.jpog.2014.01.003](https://doi.org/10.1016/j.jpog.2014.01.003).
- Oktay, K., Cil, A.P., & Bang, H. (2006). Efficiency of oocyte cryopreservation: A meta-analysis. *Fertility and Sterility*, 86, 70–80. doi: [10.1016/j.fertnstert.2006.03.017](https://doi.org/10.1016/j.fertnstert.2006.03.017).
- Okugawa, K., Kobayashi, H., Sonoda, K., Kaneki, E., Kawano, Y., Hidaka, N., ... Kato, K. (2016). Oncologic and obstetric outcomes and complications during pregnancy after fertility-sparing abdominal trachelectomy for cervical cancer: A

- retrospective review. *International Journal of Clinical Oncology*, 22, 340–346. doi: [10.1007/s10147-016-1059-9](https://doi.org/10.1007/s10147-016-1059-9).
- Pacey, A.A., & Eiser, C. (2011). Banking sperm is only the first of many decisions for men: What healthcare professionals and men need to know. *Human Fertility (Cambridge)*, 14, 208–217. doi: [10.3109/14647273.2011.634480](https://doi.org/10.3109/14647273.2011.634480).
- Parmegiani, L., Garello, C., Granello, F., Guidetti, D., Bernardi, S., Cognigni, G.E., ... Filicori, M. (2009). Long-term cryo-storage does not adversely affect the outcome of oocyte thawing cycles. *Reproductive Biomedicine Online*, 19, 374–379. doi: [10.1016/S1472-6483\(10\)60171-X](https://doi.org/10.1016/S1472-6483(10)60171-X).
- Peate, M., Meiser, B., Cheah, B.C., Saunders, C., Butow, P., Thewes, B., & Friedlander, M. (2012). Making hard choices easier: A prospective, multicentre study to assess the efficacy of a fertility-related decision aid in young women with early-stage breast cancer. *British Journal of Cancer*, 106, 1053–1061. doi: [10.1038/bjc.2012.61](https://doi.org/10.1038/bjc.2012.61).
- Peate, M., Meiser, B., Hickey, M., & Friedlander, M. (2009). The fertility-related concerns, needs and preferences of younger women with breast cancer: A systematic review. *Breast Cancer Research and Treatment*, 116, 215–223. doi: [10.1007/s10549-009-0401-6](https://doi.org/10.1007/s10549-009-0401-6).
- Pérez-Andújar, A., Newhauser, W.D., Taddei, P.J., Mahajan, A., & Howell, R.M. (2013). The predicted relative risk of premature ovarian failure for three radiotherapy modalities in a girl receiving craniospinal irradiation. *Physics in Medicine and Biology*, 58, 3107–3123. doi: [10.1088/0031-9155/58/10/3107](https://doi.org/10.1088/0031-9155/58/10/3107).
- Practice Committees of American Society for Reproductive Medicine and the Society for Assisted Reproduction. (2013a). In vitro maturation: A committee opinion. *Fertility and Sterility*, 99, 663–666. doi: [10.1016/j.fertnstert.2012.12.031](https://doi.org/10.1016/j.fertnstert.2012.12.031).
- Practice Committees of American Society for Reproductive Medicine and the Society for Assisted Reproduction. (2013b). Mature oocyte cryopreservation: A guideline. *Fertility and Sterility*, 99, 37–43. doi: [10.1016/j.fertnstert.2012.09.028](https://doi.org/10.1016/j.fertnstert.2012.09.028).
- Practice Committees of American Society for Reproductive Medicine and the Society for Assisted Reproduction. (2014). Ovarian tissue cryopreservation: A committee opinion. *Fertility and Sterility*, 101, 1237–1243. doi: [10.1016/j.fertnstert.2014.02.052](https://doi.org/10.1016/j.fertnstert.2014.02.052).
- Quinn, M.M., Cakman, H., Letourneau, J.M., Cedars, M.I., & Rosen, M.P. (2017). Response to ovarian stimulation is not impacted by a breast cancer diagnosis. *Human Reproduction*, 32, 568–574. doi: [10.1093/humrep/dew355](https://doi.org/10.1093/humrep/dew355).
- Reulen, R.C., Zeegers, M.P., Wallace, W.H., Frobisher, C., Taylor, A.J., Lancashire, E.R., ... Hawkins, M.M. (2009). Pregnancy outcomes among adult survivors of childhood cancer in the British childhood cancer survivor study. *Cancer Epidemiology, Biomarkers and Prevention*, 18, 2239–2247. doi: [10.1158/1055-9965](https://doi.org/10.1158/1055-9965).
- Roberts, J., Ronn, R., Tallon, N., & Holzer, H. (2015). Fertility preservation in reproductive age woman facing gonadotoxic treatments. *Current Oncology*, 22, 294–304. doi: [10.3747/co.22.2334](https://doi.org/10.3747/co.22.2334).
- Royal College of Physicians, The Royal College of Radiologists, Royal College of Obstetricians and Gynaecologists. (2007). The effects of cancer treatment on reproductive functions: Guidance on management. Report of a Working Party. Retrieved from https://www.rcr.ac.uk/system/files/publication/field_publication_files/Cancer_fertility_effects_Jan08.pdf.
- Ruddy, K.J., Gelber, S.I., Tamimi, R.M., Ginsburg, E.S., Schapira, L., Come, S.E., ... Partridge, A.H. (2014). Prospective study of fertility concerns and preservation strategies in young women with breast cancer. *Journal of Clinical Oncology*, 32, 1151–1156. doi: [10.1200/JCO.2013.52.8877](https://doi.org/10.1200/JCO.2013.52.8877).
- Schmidt, K.T., & Anderson, C.Y. (2012). Recommendations for fertility preservation in patients with lymphomas. *Journal of Assisted Reproduction and Genetics*, 29, 473–477. doi: [10.1007/s10815-012-9808-9](https://doi.org/10.1007/s10815-012-9808-9).
- Scottish Intercollegiate Guidelines Network. (2013). Long term follow up of survivors of childhood cancer. Retrieved from <http://www.sign.ac.uk/assets/sign132.pdf>.
- Shou, H., Chen, Y., Chen, Z., Zhu, T., & Ni, J. (2015). Laparoscopic ovarian transposition in young women with cervical squamous cell carcinoma treated by primary pelvic irradiation. *European Journal of Gynaecological Oncology*, 36, 25–29. doi: [10.12892/ejgo2607.2015](https://doi.org/10.12892/ejgo2607.2015).
- Signorello, L.B., Cohen, S.S., Bosetti, C., Stovall, M., Kasper, C.E., Weathers, R.E., ... Boice, J.D. Jr. (2006). Female survivors of childhood cancer: preterm birth and low birth weight among their children. *Journal of the National Cancer Institute*, 98, 1453–1461. doi: [10.1093/jnci/djj394](https://doi.org/10.1093/jnci/djj394).
- Signorello, L.B., Mulvihill, J.J., Green, D.M., Munro, H.M., Stovall, M., Weathers, R.E., ... Boice, J.D. Jr. (2012). Congenital anomalies in the children of cancer survivors: A report from the childhood cancer survivor study. *Journal of Clinical Oncology*, 30, 239–245. doi: [10.1200/JCO.2011.37.2938](https://doi.org/10.1200/JCO.2011.37.2938).
- Stern, C.J., Gook, D., Hale, L.G., Agresta, F., Oldham, J., Rozen, G., & Jobling, T. (2013). First reported clinical pregnancy following heterotopic grafting of cryopreserved ovarian tissue in a woman after a bilateral oophorectomy. *Human Reproduction*, 28, 2996–2999. doi: [10.1093/humrep/det360](https://doi.org/10.1093/humrep/det360).
- Tan, P.L., Wagner, J.E., Auerbach, A.D., Defor, T.E., Slungaard, A., & Macmillan, M.L. (2006). Successful engraftment without radiation after fludarabine-based regimen in Fanconi anemia patients undergoing genotypically identical donor hematopoietic cell transplantation. *Pediatric Blood and Cancer*, 46, 630–636. doi: [10.1002/pbc.20538](https://doi.org/10.1002/pbc.20538).
- Terenziani, M., Piva, L., Meazza, C., Gandola, L., Cefalo, G., & Merola, M. (2009). Oophoropexy: A relevant role in preservation of ovarian function after pelvic irradiation. *Fertility and Sterility*, 91, 15–16. doi: [10.1016/j.fertnstert.2008.09.029](https://doi.org/10.1016/j.fertnstert.2008.09.029).
- Tschudin, S., & Bitzer, J. (2009). Psychological aspects of fertility preservation in men and women affected by cancer and other life-threatening diseases. *Human Reproduction Update*, 15, 587–597. doi: [10.1093/humupd/dmp015](https://doi.org/10.1093/humupd/dmp015).
- Tschudin, S., Bunting, L., Abraham, J., Gallop-Evans, E., Fiander, A., & Boivin, J. (2010). Correlates of fertility issues in an internet survey of cancer survivors. *Journal of Psychosomatic Obstetrics and Gynaecology*, 31, 150–157. doi: [10.3109/0167482X.2010.503910](https://doi.org/10.3109/0167482X.2010.503910).
- Turan, V., Bedoschi, G., Moy, F., & Oktay, K. (2013). Safety and feasibility of performing two consecutive ovarian stimulation cycles with the use of letrozole-gonadotropin protocol for fertility preservation in breast cancer patients. *Fertility and Sterility*, 100, 1681–1685. doi: [10.1016/j.fertnstert.2013.08.030](https://doi.org/10.1016/j.fertnstert.2013.08.030).

- Van der Ven, H., Liebenthron, J., Beckmann, M., Toth, B., Korell, M., Krussel, J., ... Dittrich, R. (2016). Ninety-five orthotopic transplantations in 74 women of ovarian tissue after cytotoxic treatment in a fertility preservation network: Tissue activity, pregnancy and delivery rates. *Human Reproduction*, 31, 2031–2041. doi: [10.1093/humrep/dew165](https://doi.org/10.1093/humrep/dew165).
- van Erven, B., Gubbels, C.S., van Golde, R.J., Dunselman, G.A., Derhaag, J.G., de Wert, G., ... Rubio-Gozalbo, M.E. (2013). Fertility preservation in female classic galactosemia patients. *Orphanet Journal of Rare Diseases*, 8, 107. doi: [10.1186/1750-1172-8-107](https://doi.org/10.1186/1750-1172-8-107).
- van Wolff, M., Montag, M., Dittrich, R., Denschlag, D., Nawroth, F., & Lawrenz, B. (2011). Fertility preservation in women—a practical guide to preservation techniques and therapeutic strategies in breast cancer, Hodgkin's lymphoma and borderline ovarian tumours by the fertility preservation network FertiPROTEKT. *Archives of Gynecology and Obstetrics*, 284, 427–435. doi: [10.1007/s00404-011-1874-1](https://doi.org/10.1007/s00404-011-1874-1).
- Wallace, W.H., Smith, A., Kelsey, T., Edgar, A., & Anderson, R. (2014). Fertility preservation for girls and young women with cancer: Population-based validation of criteria for ovarian tissue cryopreservation. *The Lancet*, 15, 1129–1136. doi: [10.1016/S1470-2045\(14\)70334-1](https://doi.org/10.1016/S1470-2045(14)70334-1).
- Wennerholm, U.B., Söderström-Anttila, V., Bergh, C., Aittomäki, K., Hazekamp, J., Nygren, K.G., ... Loft, A. (2009). Children born after cryopreservation of embryos or oocytes: A systematic review of outcome data. *Human Reproduction*, 24, 2158–2172. doi: [10.1093/humrep/dep125](https://doi.org/10.1093/humrep/dep125).
- Wilkes, S., Coulson, S., Crosland, A., Rubin, G., & Stewart, J. (2010). Experience of fertility preservation among younger people diagnosed with cancer. *Human Fertility*, 13, 151–158. doi: [10.3109/14647273.2010.503359](https://doi.org/10.3109/14647273.2010.503359).
- Williams, C.L., Bunch, K.J., Stiller, C.A., Murphy, M.F., Botting, B.J., Wallace, W.H., ... Sutcliffe, A.G. (2013). Cancer risk among children born after assisted conception. *The New England Journal of Medicine*, 369, 1819–1827. doi: [10.1056/NEJMoa1301675](https://doi.org/10.1056/NEJMoa1301675).
- Willows, K., Lennox, G., & Covens, A. (2016). Fertility-sparing management in cervical cancer: Balancing oncologic outcomes with reproductive success. *Gynecologic Oncology Research and Practice*, 3, 9. doi: [10.1186/s40661-016-0030-9](https://doi.org/10.1186/s40661-016-0030-9).
- Winther, J.F., Olsen, J.H., Wu, H., Shyr, Y., Mulvihill, J.J., Stovall, M., ... Boice, J.D. Jr. (2012). Genetic disease in the children of Danish survivors of childhood and adolescent cancer. *Journal of Clinical Oncology*, 30, 27–33. doi: [10.1200/JCO.2011.35.0504](https://doi.org/10.1200/JCO.2011.35.0504).
- Wo, J.Y., & Viswanathan, A.N. (2009). Impact of radiotherapy on fertility, pregnancy, and neonatal outcomes in female cancer patients. *International Journal of Radiation Oncology Biology Physics*, 73, 1304–1312. doi: [10.1016/j.ijrobp.2008.12.016](https://doi.org/10.1016/j.ijrobp.2008.12.016).
- Woodruff, T.K., Clayman, M.L., & Waimey, K.E. (2014). *Oncofertility communication: Sharing information and building relationships across disciplines*. New York, NY: Springer.
- Yoon, M. (2012). Development of and current status of proton therapy for lung cancer in Korea. *Thoracic Cancer*, 3, 1–7. doi: [10.1111/j.1759-714.2011.00085.x](https://doi.org/10.1111/j.1759-714.2011.00085.x).
- Youssef, M.A., Van der Veen, F., Al-Inany, H.G., Griesinger, G., Mochtar, M.H., Aboulfoutouh, I., & van Wely, M. (2014). Gonadotropin-releasing hormone agonist versus HCG for oocyte triggering in antagonist assisted reproductive technology cycles. *Cochrane Database Systematic Review*, CD008046. doi: [10.1002/14651858.CD008046.pub4](https://doi.org/10.1002/14651858.CD008046.pub4).